

# Chlorophyll Fluorescence Analysis of Cyanobacterial Photosynthesis and Acclimation

DOUGLAS CAMPBELL,<sup>1\*</sup> VAUGHAN HURRY,<sup>2</sup> ADRIAN K. CLARKE,<sup>2</sup> PETTER GUSTAFSSON,<sup>2</sup>  
AND GUNNAR ÖQUIST<sup>2</sup>

*Department of Biology, Mount Allison University, Sackville, New Brunswick E4L 1G7, Canada,<sup>1</sup> and  
Department of Plant Physiology, University of Umeå, S901-87 Umeå, Sweden<sup>2</sup>*

INTRODUCTION .....	667
Principles of Modulated Fluorescence Analysis .....	667
Goals and Scope.....	668
INTERPRETING CYANOBACTERIAL FLUORESCENCE SIGNALS.....	668
A Distinct Photosynthetic System Yields Distinct Fluorescence Signals.....	668
$F_o$ , $F_v/F_m$ , and $F_o'$ in Cyanobacteria versus Plants .....	670
PHOTOCHEMICAL QUENCHING AND EXCITATION PRESSURE.....	672
PREDICTING LIGHT ACCLIMATION STATUS.....	673
State Transitions Dominate Nonphotochemical Quenching of Cyanobacterial Fluorescence.....	673
Predicting the Acclimated Light Intensity from $q_N$ Light Response .....	675
Inorganic Carbon Accumulation and Fluorescence Quenching.....	676
State Transitions Can Be Measured Nondestructively .....	676
PREDICTING PHOTOSYNTHESIS FROM FLUORESCENCE .....	676
APPLYING FLUORESCENCE ANALYSIS TO DIFFERENT CYANOBACTERIA .....	677
CONCLUSIONS AND PROSPECTS.....	678
Tracking Acclimation Status in the Laboratory and Field.....	678
Conclusions.....	678
APPENDIX .....	679
Cyanobacterial Cultures and Pigment Measurements.....	679
Modulated Fluorometer Configuration and Measurement Procedures .....	679
77K Fluorescence Emission Spectra .....	680
Alternate Fluorescence Quenching Parameters.....	680
ACKNOWLEDGMENTS .....	680
REFERENCES .....	680

## INTRODUCTION

### Principles of Modulated Fluorescence Analysis

Chlorophyll fluorescence analysis allows noninvasive, near-instantaneous measurement of key aspects of photosynthetic light capture and electron transport. For natural samples, fluorescence signals are specific to photobionts and allow in situ measurements of small (61) or dilute (65, 130) mixed natural populations. For molecular studies, fluorescence signals can be used for rapid screening of mutant or transgenic colonies and cultures and for tracking physiological processes during gene regulation experiments. Rapid screening has become increasingly important with the advent of genomic sequencing and saturation mutagenesis. Therefore, applications of chlorophyll fluorescence are expanding in both field and laboratory settings.

In cyanobacteria, the photosynthetic system is tightly connected to the other principal metabolic paths and is in itself a major metabolic sink for iron, nitrogen, and carbon skeletons. Therefore, chlorophyll fluorescence signals can provide rapid, real-time information on both photosynthesis and the overall acclimation status of cyanobacteria. We and other groups have

been adapting to cyanobacteria techniques of in vivo fluorescence analysis originally developed for plants (5, 20–28, 58, 61, 69, 74, 80, 87–91, 112, 116, 128, 133, 138, 141, 142, 150).

Fluorescence analysis depends on the phenomenon that when a pigment absorbs the energy of a photon and enters an excited electronic state, there are essentially four routes for the return to ground state: (i) photochemical reactions in which the excited electron leaves the pigment molecule and enters an electron transport chain, as occur in specific chlorophylls in photosynthetic reaction centers; (ii) heat dissipation, in which the excited electron returns to ground state by releasing heat; (iii) transfer of the excitation energy to an adjacent pigment, as occurs in the light-harvesting antenna systems of photosynthetic organisms; and (iv) emission of a fluorescence photon, of a wavelength longer than that of the photon initially absorbed. These four processes are in competition, and for a given excited molecule, the path with the largest first-order rate constant predominates. For biological systems, the overall chlorophyll fluorescence yield is usually low, and in vivo chlorophyll fluorescence from photosystem II (PS II) predominates (38, 66, 112). In cyanobacteria, phycobiliproteins also contribute fluorescence, which overlaps with the spectrum of chlorophyll emission.

Although PS II fluorescence is a minor pathway for excitation dissipation, it competes with the quantitatively more important energy dissipation routes of PS II photochemistry, exciton transfer to other pigment systems (such as PS I), and heat dissipation. Therefore, changes in photochemistry or in

\* Corresponding author. Mailing address: Department of Biology, Mount Allison University, Sackville, New Brunswick E4L 1G7, Canada. Phone: (506) 364 2521. Fax: (506) 364 2505. E-mail: dcampbell@mta.ca.

TABLE 1. Fluorescence levels and associated light treatments used for cyanobacterial fluorescence quenching analysis<sup>a</sup>

Fluorescence level	Light treatment (weak modulated measuring beam throughout)
$F_O$	Dark
$F_{M\text{dark}'}$	Dark + saturating flash
$F_S$	Actinic light
$F_O'$	Dark or weak far-red
$F_M'$	Actinic light + saturating flash
$F_M$	Actinic light + DCMU

<sup>a</sup> For further details see Appendix.

the two nonphotochemical routes (energy transfer and heat emission) cause changes in the fluorescence yield from PS II (13, 66, 112). When the potentials for photochemistry and nonphotochemical dissipation are minimal, the fluorescence yield is maximal. Quenching or lowering of the fluorescence yield below its maximum occurs when excitation flow increases to the competing photochemical or nonphotochemical pathways.

To deduce information on photosynthesis from analysis of fluorescence quenching, one assumes that changes in fluorescence yield reflect proportional changes in the competing de-excitation pathways of photochemistry, exciton transfer, and heat dissipation. This basic assumption is not strictly valid (53, 56, 144). Nevertheless, the fluorescence signal is rich in information, and in plants the parameters  $F_V/F_M$ ,  $F_V'/F_M'$ ,  $q_P$ ,  $q_N$ , NPQ, and  $\phi\text{PS II}$  are empirically verifiable indices of photosynthetic performance and acclimation status (12, 36, 45, 66, 67, 82, 99, 102–104, 106, 107, 127, 129, 134, 145, 153).

For any pigment, the level of fluorescence emission depends on the pigment concentration, the excitation light intensity, and the fluorescence yield or efficiency of fluorescence emission. For fluorescence quenching analysis, the excitation intensity and pigment concentration must be constant, so that changes in fluorescence reflect the changes in fluorescence yield which result from the competing photochemical and nonphotochemical deexcitation pathways. Changes in pigment concentration are generally not a concern over the brief periods of fluorescence measurements.

Modulated fluorometers are currently widely used to measure in vivo chlorophyll fluorescence from plants and increasingly from cyanobacteria in both the laboratory and the field (128, 130). This review concentrates on data obtained with modulated fluorometers, although other approaches are also used (40). Modulated fluorometers specifically detect and amplify only the fluorescence excited by a weak, constant measuring beam consisting of a train of low light pulses at a frequency of 1 to 100 kHz. Therefore, the excitation intensity is constant and changes in the fluorometer signal reflect changes in fluorescence yield. The modulated measuring beam is sufficiently weak that it drives essentially no photosynthesis, allowing determination of the fluorescence yield of dark-adapted samples. Furthermore, since the detection system ignores fluorescence excited by other light, it is possible to change the actinic light and provide multiple saturating pulses of light over the course of one measurement (Table 1; see Fig. 2). The fluorescence yield can therefore be measured under different levels of actinic light, and saturating flashes can be used at any point to momentarily close all PS II centers and drive photochemical quenching to zero (see Fig. 2).

## Goals and Scope

In this review, we discuss pulse-amplitude modulated fluorescence as a rapid, noninvasive monitor of acclimation and photosynthesis in cyanobacteria and cyanolichens. We do not cover the biophysical mechanisms underlying chlorophyll fluorescence emission, which are well reviewed elsewhere (31, 38, 66, 112, 127, 137, 141, 144). Rather, we summarize some of the potentials and limitations of fluorescence analysis for extracting physiologically and ecologically useful information from cyanobacteria, whose photosynthetic physiology (see Fig. 1) and fluorescence patterns (see Fig. 2) differ in important respects from those of plants (20, 22, 91, 112, 128, 141). In particular, we demonstrate how characteristic changes in non-photochemical quenching of fluorescence can be used to estimate the light level to which the sample is acclimated. This information can then be used in conjunction with the  $\phi\text{PS II}$  parameter to estimate electron transport under acclimated conditions.

## INTERPRETING CYANOBACTERIAL FLUORESCENCE SIGNALS

### A Distinct Photosynthetic System Yields Distinct Fluorescence Signals

The central PS II and PS I photosynthetic complexes are very similar in plants and cyanobacteria, as are many elements of the light capture, electron transport, and carbon dioxide fixation systems. Nevertheless, cyanobacteria are metabolically flexible prokaryotic organisms, with several key structural and metabolic distinctions which strongly influence the nature and interpretation of their fluorescence signals (Fig. 1).

In cyanobacteria, the principal light-harvesting complexes are phycobilisomes peripheral to the thylakoid membranes, rather than the integral membrane chlorophyll-*a/b* binding proteins which capture light in plants. Cyanobacterial phycobilisomes diffuse along the surface of the thylakoids, at a rate sufficient to allow movement from PS II to PS I within 100 ms (96). This distinction in light capture structures between plants and cyanobacteria has many metabolic and functional consequences (7). In particular, the cellular phycobiliprotein content influences cellular fluorescence yield. Furthermore, cyanobacteria have high and variable ratios between PS I and PS II complexes (98, 112), so that in comparison with plants, PS II accounts for relatively little of the cellular chlorophyll. This can also influence the interpretation of fluorescence signals, since the variable fluorescence component arises from PS II while the constant or  $F_0$  fluorescence component contains emissions from PS II, phycobiliproteins, and possibly also PS I chlorophyll (112).

Photosynthetic and respiratory electron flow both occur in cyanobacterial thylakoid membranes (62, 121), sometimes simultaneously, and they share numerous electron transport intermediates (Fig. 1). Under illumination, there is net input of electrons into the transport system from the water-splitting activity of PS II. Under light or dark conditions, there are variable electron fluxes from NAD(P)H, which is oxidized by one or more thylakoid-bound dehydrogenases (11, 54, 85, 86, 121, 139). Electrons from ferredoxin can also enter the transport system, possibly passing via the same complex(es) that catalyze NAD(P)H oxidation (86). In a photoautotroph, electrons derived from NAD(P)H or ferredoxin are clearly not a net input of reductant into the system; rather, they represent some form of cyclic flow, since the electrons used to reduce NAD(P)H or ferredoxin derive ultimately from the water-




FIG. 1. Cyanobacterial thylakoid electron transport. This schematic diagram is based on data primarily from *Synechocystis* sp. strain PCC 6701, which contains phycoerythrin, phycocyanin, and allophycocyanin pigment proteins in the phycobilisomes. The phycobilisomes move rapidly along the surface of the thylakoid membranes (96), so the phycobilisome-PS II dimer complex is transient. Excitation absorbed by the phycobilisome can reach either PS II or PS I, particularly in cells in state II. This excitation flow may involve specialized subunits of the phycobilisome core (not shown here). The composition and organization of the phycobilisome rods and core is variable in different cyanobacteria; the three-cylinder core and six-peripheral rod configuration is common, but in *Synechococcus* sp. strain PCC 7942 the core contains only two cylinders. The PS I-PS II stoichiometry is usually higher than 1:1; in the *Synechococcus* cells used for most experiments described in later figures, the ratio was 2 to 3 PS I/PS II. There are multiple interacting and flexible paths of electron flow including linear flow from water to NADPH; several possible cyclic pathways around subsections of the transport system; pseudocyclic flows from water with electron donation back to oxygen; and respiratory flows of electrons derived from reserve molecules. Some possible flows are indicated by black arrows. The donor-acceptor stoichiometries of electron transfers are not shown, but various redox centres carry different numbers of electrons, from 1 (e.g., PC, cytochrome *c*<sub>553</sub>, Fd, and Flvd), 2 [e.g., PQ and NAD(P)H] and even 4 (the Mn complex of PS II). The redox reactions are reversible depending upon the oxidation-reduction status of the acceptor-donor pair and the local proton concentration, so that in some cases the indicated direction of electron flow could be reversed. Proton uptake and transmembrane transport are indicated by dashed arrows; other proton translocation pathways may also exist. The plastoquinone-plastoquinol pool can be reduced by electrons from PS II, from an NAD(P)H dehydrogenase(s) (NDH) whose composition and substrate specificities vary between strains, and from the "r" site of the cytochrome *bf* complex. Plastoquinone reduction by NDH is the entry point for electrons derived from respiration. The various plastoquinone reductions involve proton uptake from the stroma, while oxidation of a plastoquinol at the "o" site of the cytochrome *bf* complex releases two protons to the lumen. As indicated, electron and ATP flow to carbohydrate and nitrogen metabolism can have strong and rapid effects on thylakoid function, while electrons derived from carbohydrate reserves also enter the thylakoid system and influence photosynthetic function. During ATP synthesis, protons enter a channel from the lumen formed by the a subunit of CF<sub>0</sub>, and their exit to the cytosol is coupled to rotation of the ring of c subunits through directed diffusion (37). The c-ring rotation drives rotation of the  $\gamma$  subunit of CF<sub>1</sub> within the  $\alpha_3\beta_3$  ring (117), which in turn drives a sequence of conformational changes in three identical ATP/ADP binding sites. The changes in binding site lead to phosphorylation of ADP and expulsion of ATP from the site. The stoichiometry is 1 ATP/4H<sup>+</sup> passing through the complex (146). Abbreviations: hv, photons of visible light; PE, phycoerythrin  $\alpha_3\beta_3$  trimers; CPC, phycocyanin  $\alpha_3\beta_3$  trimers; AP, allophycocyanin rods of the phycobilisome core, composed partly of  $\alpha_3\beta_3$  trimer disks along with other related phycobilin-binding proteins; D1 and D2, core polypeptide dimer of PS II which binds the redox cofactors; Cyt b<sub>559</sub>, cytochrome *b*<sub>559</sub> in the PS II core; Mn<sub>4</sub>, manganese cluster of the oxygen evolving complex; 9 and 33, 9- and 33-kDa subunits of the oxygen evolving complex of PS II; CP43 and CP47, 43- and 47-kDa chlorophyll protein complexes associated with the PS II core; P<sub>680</sub>, dimeric chlorophyll center which is photooxidized in PS II; Pheo, pheophytin primary electron acceptor of PS II; Q<sub>A</sub>, the quinone secondary electron acceptor of PS II; Q<sub>B</sub>, a plastoquinone bound to PS II which accepts two electrons from Q<sub>A</sub> and equilibrates with the thylakoidal plastoquinone-plastoquinol pool; NDH, NAD(P)H dehydrogenase (in various strains there are different forms of the complex with differing activities and specificities for NADH or NADPH); Cyt b<sub>6</sub>, a cytochrome containing both low- and high-potential heme centers which are involved in a Q-cycle electron flow from plastoquinol bound to the "o" site to plastoquinone bound to the "r" site (this cycle results in proton translocation); S<sub>IV</sub>, subunit IV of the cytochrome *bf* complex; FeS, an iron-sulfur redox center; PC, plastocyanin, a copper-containing luminal single-electron transport protein; Cyt c<sub>553</sub>, cytochrome *c*<sub>553</sub>, a heme-containing luminal single-electron transport protein (plastocyanin and cytochrome *c*<sub>553</sub> can be reciprocally regulated in response to copper and iron availability); PsaA and PsaB, related chlorophyll binding proteins which form the core of PS I; P<sub>700</sub>, the chlorophyll which is photooxidized in PS I; A<sub>0</sub>, A<sub>1</sub>, F<sub>X</sub>, F<sub>A</sub>, F<sub>B</sub>, bound redox intermediates of PS I; Flvd, flavodoxin (a flavin protein which is a cytosolic mobile single electron carrier that can accept electrons from PS I and that can substitute for ferredoxin, particularly under low-iron conditions); Fd, ferredoxin (an iron protein which is a cytosolic mobile single-electron carrier that can accept electrons from PS I and can transfer the electrons to NADPH or participate directly in some biosynthetic reactions, particularly in nitrogen metabolism); FNR, ferredoxin/flavodoxin NADPH oxidoreductase; Cyt ox, the cytochrome oxidase complex involved in respiratory electron transport (it can also withdraw electrons from photosynthetic electron transport, particularly under excess light);  $\alpha$  to  $\gamma$ , subunits of the CF<sub>1</sub> complex of ATP synthase; a to c, subunits of the CF<sub>0</sub> complex of ATP synthase. Modified and redrawn from reference 19 with permission of the publisher.

splitting activity of PS II. Typically, cyclic electron flow is used to describe flow from PS I via ferredoxin and/or NADPH (54, 86) directly back to the intersystem transport chain. This cyclic flow can drive proton translocation through localized reductions of plastoquinone at the cytosolic side of the thylakoid, with concomitant proton uptake, and plastoquinol oxidation

near the luminal side, with proton release to the lumen (55) (Fig. 1). More generally, electrons derived from the oxidation of carbohydrates or other reserve molecules can be carried by NAD(P)H into the thylakoid intersystem transport chain via the thylakoid-bound dehydrogenase(s). These electrons from the reserve molecules are derived ultimately from PS II water

splitting and, upon reentry to the thylakoid system, can pass to oxygen under light or dark or to PS I (139) under illumination. The reserves thus act as an electron bank so that the flow into the thylakoid system can be offset in time from the original photosynthetic production of reductant, with important regulatory consequences (34, 84, 94).

In all known cases, electrons from these various inputs come together at the cytochrome *bf* complex, which is a plastoquinol oxidoreductase (68) (Fig. 1). There are two plastoquinone binding sites in the complex, which allow for a Q cycle, in which some of the electrons removed from plastoquinol at the luminal side of the membrane are cycled within the complex and passed back to plastoquinone bound near the cytoplasmic side of the membrane. This branch of the transport chain allows the cyanobacteria to increase the number of protons translocated per net electron passing through the transport chain. Since the reduction involves proton uptake from the cytosol and the oxidation releases protons to the lumen, electron flux through the Q cycle must respond sensitively to the magnitude of the proton gradient across the membrane.

The primary electron flux through the cytochrome *bf* complex is from plastoquinone to luminal electron carriers, primarily plastocyanin or cytochrome  $c_{553}$  (81, 113), which transport the electrons either to PS I or to a cytochrome oxidase complex which may include cytochrome *c*(m) (81). Plastocyanin and cytochrome  $c_{553}$  can each fulfill transport roles to PS I or the cytochrome oxidase (81). Double-inactivation mutants mutated in both proteins are inviable in some (81) but not all (155) strains, so that some strains must have an alternate route for electron flow away from cytochrome *bf*. Although single-inactivation mutants mutated in one or the other protein are viable (26, 71, 81), the loss of one protein can lower the capacity for electron flux away from PS II, particularly under conditions of excess excitation (26). Conversely, overexpression of heterologous plastocyanin in *Synechococcus* can increase the electron transport capacity (44). Thus, although partially complementary, the two proteins may play somewhat distinct functional roles. Furthermore, plastocyanin contains a copper redox cofactor while cytochrome  $c_{553}$  contains an iron redox cofactor, and in some strains they are differentially regulated in response to copper and iron availability (15, 16, 119, 120, 154). Another example of alternate electron carriers is the iron-sulfur protein ferredoxin, which accepts electrons from PS I but which can be replaced under conditions of iron stress by flavodoxin (70, 75).

The flow of electrons to oxygen as a final acceptor responds rapidly to environmental and metabolic conditions and can be an important element in preventing overreduction of PS II and the intersystem transport chain under excess illumination (22a, 88, 147). This flow to oxygen can be mediated by cytochrome oxidase activity (131) or by photoreduction of oxygen by electrons from PS I, either directly (3, 79) or via ferredoxin (43, 50).

Carbon metabolism and nitrogen metabolism in the prokaryotic cyanobacteria occur in close proximity to the cytosolic surface of the thylakoids and so have strong and direct influences on electron transport and hence on fluorescence (63, 83, 84, 92, 116, 122), both as sinks for ATP and electrons and as sources of electrons extracted from reserve molecules.

In summary, this system forms a web of electron sources and sinks, linked by interconnected redox intermediates, that allows for flexible and rapid shifts in electron fluxes in response to environmental or metabolic changes (5, 54, 55, 74, 80, 81, 85, 86, 131, 135, 147). Furthermore, through shared electron transport carriers, respiration directly influences the photosynthetic regulatory status and vice versa (34, 84, 94). Several of the

characteristic properties of cyanobacterial fluorescence signals result from these respiration/photosynthesis interactions, including their distinct patterns of photochemical and nonphotochemical quenching.

In plants, a cycle of conversions of xanthophyll carotenoids is driven by the *trans*-thylakoid  $\Delta$ pH gradient and is implicated in regulating nonphotochemical dissipation of excess light energy (1, 46, 47, 57). Cyanobacteria lack this cycle (33) but have alternate strategies to cope with excess excitation (105, 147). Finally, cyanobacteria show changes in the functional organization of the light capture system, termed state transitions, which can result in large changes in the PS II fluorescence yield depending upon the level of illumination (7, 14, 97, 111, 112). In contrast, in higher plants, state transitions have relatively minor influences on PS II fluorescence (66). This review deals now with how the distinct organization and function of cyanobacterial photosynthesis lead to opportunities and limitations for chlorophyll fluorescence analysis of cyanobacteria.

### $F_O$ , $F_V/F_M$ , and $F_O'$ in Cyanobacteria versus Plants

Ting and Owens (141) have shown that for any chlorophyll-containing suspension, the values of  $F_O$ ,  $F_M$  and  $F_V/F_M$  measured with a modulated fluorometer vary somewhat with pigment concentration. Therefore, within a set of experiments, the chlorophyll concentration should be standardized if precision is required or, alternately, a small correction could be introduced to compensate for variation in the chlorophyll concentration. In our experiments, chlorophyll concentrations from 1 to 3  $\mu$ g/ml gave results sufficiently consistent for quenching analysis. The use of alternate cuvettes and detectors can greatly extend this concentration range (125).

A more fundamental problem with the measurement of  $F_O$  in cyanobacteria is that  $F_O$  fluorescence varies considerably depending on the cellular phycobiliprotein concentration. Figure 3 illustrates that as the phycocyanin/chlorophyll ratio of wild-type *Synechococcus* sp. strain PCC 7942 rises,  $F_O$  fluorescence also increases, particularly once the phycocyanin content is increased above a threshold level. This phycobiliprotein contribution to  $F_O$  fluorescence is not influenced by changes in the redox state of PS II (51, 72, 111). It could be a low-yield fluorescence emission from coupled phycobilisomes or a high-yield emission from a small population of uncoupled phycobilisomes (96) or free phycobiliproteins. The exact source(s) of this phycobiliprotein fluorescence could be addressed by studies using cyanobacteria with a range of phycobiliprotein contents analyzed with the range of different color light-emitting diodes now available as modulated fluorescence excitation sources (65). In a *Synechococcus* sp. strain PCC 7942 mutant lacking phycocyanin,  $F_O$  is low (Fig. 3), which confirms the  $F_O$ /phycobiliprotein correlation observed in the wild type. Furthermore, in this mutant strain,  $F_V/F_M$  under acclimated growth is about 0.75 (157), as opposed to values of 0.4 to 0.6, typical of wild-type *Synechococcus* grown under the same conditions.

In plants  $F_V/F_M [(F_M - F_O)/F_M]$  (Fig. 2) is well verified as an index of the maximal photochemical efficiency of PS II (12), but this interpretation depends on both  $F_O$  and  $F_V$  originating predominantly from PS II. This assumption is not valid for cyanobacteria (20, 111, 112, 128), since phycobiliprotein fluorescence contributes to  $F_O$  and PS II accounts for only a small proportion of total chlorophyll. In higher plants under ideal conditions,  $F_V/F_M$  is near 0.8 and lower values reflect inhibition of PS II function (12). In cyanobacteria, changes in  $F_V/F_M$  under conditions of constant pigment content correlate well with changes in independent measurements of PS II function




FIG. 2. Fluorescence emission trace for cyanobacterial quenching analysis. This trace from *Synechococcus* sp. strain PCC 7942 shows a typical cyanobacterial response over a series of increasing light intensities. The brief pulses of saturating light result in a rapid increase in fluorescence as PS II centers close transiently. The measurement terminates with addition of DCMU, which closes PS II centers, causing a rapid rise in fluorescence followed by a slower fluorescence rise phase as the cells go to full state I. Modified from reference 23 with permission of the publisher.

such as oxygen evolution (24, 27, 28, 80), but the absolute level of  $F_V/F_M$  is not a reliable indicator of PS II function. The distortion of  $F_O$  fluorescence is pronounced only at high cellular concentrations of phycocyanin (Fig. 3), which may be achieved primarily under nutrient-rich artificial culture conditions, as used in our experiments to date. When interpreted with caution,  $F_V/F_M$  is still a useful parameter, particularly if the same sample is monitored repeatedly over time and if the cellular pigment content is constant. As expected, the prochlorophyte *Prochloron* shows a high value of  $F_V/F_M$  (126), consis-




FIG. 3.  $F_O$  fluorescence increases with the phycocyanin content in *Synechococcus* sp. strain PCC 7942.  $F_O$  fluorescence is normalized to the molar chlorophyll concentration to allow for different culture concentrations and plotted against the molar ratio of phycocyanin (PC) to chlorophyll (Chl). Each point is a single determination on an independently grown culture.  $\circ$ , wild-type cells;  $\blacksquare$ , mutant strain which lacks phycobilisome rods and contains no phycocyanin. Note that at low phycocyanin contents,  $F_O$  in the wild type falls toward the level in the rodless mutant. Modified from reference 23 with permission of the publisher.

tent with the lack of phycobilisomes and the chlorophyll-based antenna system in this group of cyanobacterial relatives.

$F_O'$  is the minimal fluorescence level with all PS II reaction centers open. It is measured with cells under a given light acclimation status but transferred briefly to darkness or far-red light. The determination of  $F_O'$  is a problematic aspect of quenching analysis in cyanobacteria, since under moderate light intensities  $F_O'$  is often very close to the steady-state  $F_S$  fluorescence level (Fig. 2). Furthermore, unlike in plants,  $F_O'$  in cyanobacteria is usually higher than the  $F_O$  fluorescence as a result of the dark-to-light increase in PS II fluorescence yield, i.e., the state transition (see below). It is therefore difficult to distinguish the initial small but rapid drop in fluorescence yield as PS II centers open, from the slower state transition-dependent decline to the dark  $F_O$  fluorescence level. Computerized data logging might alleviate this problem by resolving the fluorescence relaxation kinetic phases.

In higher plants,  $F_V'/F_M'$ , defined as  $(F_M' - F_O')/F_M'$  (Fig. 2), reflects the photochemical efficiency of open PS II centers under a given light acclimation status (45).  $F_V'/F_M'$  generally varies inversely with  $q_N$  (see below), since nonphotochemical energy dissipation lowers the photochemical efficiency of PS II below the maximum levels reflected by  $F_V/F_M$ . A drop in  $F_V'/F_M'$ , as occurs during photoinhibition of PS II activity, also feeds through and results in a drop in  $F_V'/F_M'$ . Thus, in a plant, changes in the  $F_V'/F_M'$  parameter reflect the combined regulation of PS II through both reversible nonphotochemical quenching and photoinhibitory inactivation of PS II.

In cyanobacteria, changes in  $F_V'/F_M'$  also combine nonphotochemical influences on PS II function and photoinhibitory inactivation of PS II. As described below, nonphotochemical quenching of cyanobacterial PS II fluorescence results primarily from changes in excitation distribution between the two photosystems rather than from excitation dissipation. Therefore, in a cyanobacterium, a drop in  $F_V'/F_M'$  can result from a photoinhibitory drop in  $F_V/F_M$  or from a regulatory redistribu-




FIG. 4. Photoinhibition of oxygen evolution correlates with declines in  $F_V'/F_M'$  in *Synechococcus* sp. strain PCC 7942. Cultures were subjected to a photoinhibitory decrease in growth temperature from 37 to 25°C. Oxygen evolution and  $F_V'/F_M'$  were monitored and expressed as a percentages of the pretreatment control values. Four cultures were used.  $y = 0.79x + 21$ ;  $R^2 = 0.89$ .

tion of excitation from PS II to PS I. In plant fluorescence analysis, a common implicit assumption is that down-regulation of PS II reflects overall down-regulation of photosynthetic electron transport. This assumption is not applicable to cyanobacteria, which have more flexible excitation distribution and electron transport systems. Furthermore, cyanobacterial  $F_V'/F_M'$  suffers the same limitations as described above for  $F_V/F_M$ , which are further compounded by the difficulty of measuring the  $F_O'$  fluorescence level. Nevertheless, as shown in Fig. 4, changes in measured  $F_V'/F_M'$  correlate well with changes in oxygen evolution during a photoinhibitory treatment (24, 27). Cyanobacterial  $F_V'/F_M'$  is a useful integrated measure of PS II activity, even though various mechanisms may underlie the changes in PS II function.

#### PHOTOCHEMICAL QUENCHING AND EXCITATION PRESSURE

Photochemical and nonphotochemical quenching measure changes in variable fluorescence, which derives from PS II; they are therefore less susceptible to distortion from non-PS II contributions to  $F_O$  fluorescence. They involve minimal mechanistic assumptions (145), although the terms  $q_P$  and  $q_N$  often carry mechanistic associations which are not applicable to cyanobacteria.

Photochemical quenching reflects a lowering of fluorescence below maximal levels through photochemical competition with fluorescence emission. Thus, when all PS II reaction centers are open and the potential for photochemistry is maximal, photochemical quenching of fluorescence is also maximal and fluorescence yield is low ( $F_O$  or  $F_O'$ ). Conversely, when all PS II centers are closed and no photochemistry can occur, photochemical quenching is zero and the fluorescence yield is maximal ( $F_M$  or  $F_M'$ ). In practice, photochemical quenching is quantified by the photochemical quenching coefficient (145):

$$q_P = (F_M' - F_S)/(F_M' - F_O')$$




FIG. 5. Photochemical quenching of fluorescence plotted against light intensity. Typical light response curves of photochemical quenching in a cyanobacterium, *Synechococcus* sp. strain PCC 7942 (solid line), and in rye for a plant-type pattern (dotted line) are shown. For comparison, light intensity is expressed relative to the growth light intensity; 1 = growth level. The actual growth light intensities were 50  $\mu\text{mol}$  of photons  $\text{m}^{-2} \text{s}^{-1}$  for the cyanobacterium and 250  $\mu\text{mol}$  of photons  $\text{m}^{-2} \text{s}^{-1}$  for the rye.

This parameter gives the position of steady-state fluorescence,  $F_S$ , on the scale from  $F_O'$  (all PS II centers open) to  $F_M'$  (all PS II centers closed) (Fig. 2). Thus, if steady-state fluorescence is equal to  $F_O'$ ,  $q_P = 1$ , indicating that all PS II centers are open. If steady-state fluorescence equals  $F_M'$ ,  $q_P = 0$ , indicating that all PS II centers are closed. Between these extremes, progressive reaction center closure is reflected by a declining  $q_P$ , although the relationship is not strictly linear (20) because of excitation migration from closed to open reaction centers (66) and fluorescence quenching by oxidized plastoquinone (148). The absolute level of  $F_O$  or  $F_O'$  does not distort the calculation, since it is scaled to variable fluorescence,  $F_V' = F_M' - F_O'$ , under a given condition.

$q_P$  reflects the balance between excitation of PS II centers, which closes them, and removal of electrons from PS II by the electron transport chain, which reopens the centers. This balance, or excitation pressure on PS II, responds not only to incident light intensity (27) but also to factors influencing electron flow away from PS II, such as temperature (24, 59, 82, 101, 102, 106, 107) and the availability of terminal electron acceptors such as  $\text{CO}_2$  or  $\text{O}_2$  (88, 147). Indeed, the pivotal position of PS II in photosynthetic electron transport means that environmental and metabolic signals are integrated into  $q_P$ , which is thus a general index of the balance between energy capture and consumption.

As shown in Fig. 5, in cyanobacteria and cyanolichens,  $q_P$  typically stays high over a broad range of incident light intensity, up to 10 times higher than the growth light intensity (21, 23, 27, 28, 80, 138). This contrasts sharply with the pattern typical of higher plants, where  $q_P$  falls progressively as the light intensity exceeds the growth light. This cyanobacterial capacity to maintain PS II centers open under excess light reflects a complex and flexible electron transport system (5, 44, 55, 85, 86, 131, 135), as well as a generally high PS I/PS II ratio (23, 42, 98, 112). In particular, cyanobacteria have a very high and flexible capacity to remove electrons from PS II, with oxygen as a terminal acceptor for electron flow from water (21a, 88, 91, 128, 147). This flow is low under low light, variable but significant at the growth light intensity, and large under excess light (see below). It thus serves to buffer PS II from excess excitation by removing electrons as required. At least part of this flow is




FIG. 6. Photochemical quenching of fluorescence under DCMU treatment compared with oxygen evolution in *Synechococcus* sp. strain PCC 7942. DCMU inhibits oxygen evolution (○), closing PS II centers in parallel ( $q_p$ ) (■). Results are means and standard errors of measurements on the same culture at 37°C ( $n = 3$ ). Modified from reference 24 with permission of the publisher.

sensitive to cyanide inhibition, suggesting a contribution from respiratory electron flow through cytochrome oxidase (131).

In spite of the different patterns of  $q_p$  in cyanobacteria and plants, the parameter does successfully measure PS II closure in cyanobacteria, as shown by titration with 3-(3,4-dichlorophenyl)-1,1-dimethylurea (DCMU). This inhibitor binds to the  $Q_B$  binding site of PS II, blocking electron flow and causing reaction center closure. Figure 6 shows that in *Synechococcus* sp. strain PCC 7942, as expected,  $q_p$  and oxygen evolution drop in parallel upon progressive DCMU inhibition of PS II. Furthermore, our work on *Synechococcus* sp. strain PCC 7942 showed that although this strain has a strong capacity to maintain PS II open, even fractional closure of the reaction centers can lead to photoinhibition and large changes in gene expression, indicative of active acclimation processes (24, 27, 28) in response to relatively small drops in the proportion of open reaction centers.

**PREDICTING LIGHT ACCLIMATION STATUS**

**State Transitions Dominate Nonphotochemical Quenching of Cyanobacterial Fluorescence**

Nonphotochemical quenching reflects any process other than photochemistry which lowers the yield of variable fluorescence. It can be quantified by using the coefficient  $q_N$  (145):

$$q_N = 1 - [(F_M' - F_O') / (F_M - F_O)] = 1 - (F_V' / F_V)$$

$q_N$  compares the span of variable fluorescence under a given condition,  $F_V' = (F_M - F_O')$ , with the maximum potential variable fluorescence,  $F_V = (F_M - F_O)$  (Fig. 2). In cyanobacteria, variable fluorescence appears to arise essentially from PS II, while as discussed, the  $F_O'$  and  $F_O$  fluorescence levels arise only partly from PS II. The cyanobacterial  $F_O'$  and  $F_O$  signals detected by a modulated fluorometer with a red-modulated light-emitting diode (LED) each contain a contribution from phycocyanin fluorescence, which we believe is fairly constant over the course of a measurement. This underlying background fluorescence is subtracted out as an equivalent component of both  $F_O'$  and  $F_O$  and so does not seriously distort the calculation of  $q_N$ .

Significant drops in  $F_O$  fluorescence during illumination can




FIG. 7. Nonphotochemical quenching of fluorescence plotted against light. Typical light response curves of nonphotochemical quenching in a cyanobacterium, *Synechococcus* sp. strain PCC 7942 (solid line), and in rye for a plant-type pattern (dotted line) are shown. For comparison, light intensity is expressed relative to the growth light intensity; 1 = growth level. The actual growth light intensities were 50  $\mu\text{mol}$  of photons  $\text{m}^{-2} \text{s}^{-1}$  for the cyanobacterium and 250  $\mu\text{mol}$  of photons  $\text{m}^{-2} \text{s}^{-1}$  for the rye.

distort  $q_N$  in higher plants. This  $F_O$  quenching is quantified as follows (9):

$$1 - q_O = F_O' / F_O$$

A modified calculation of the  $q_N$  coefficient compensates for  $q_O$  (9):

$$q_N = 1 - \{F_V' / [F_V (F_O' / F_O)]\}$$

In cyanobacteria,  $F_O'$  is usually close to or somewhat higher than  $F_O$  ('negative'  $q_O$ ), and both  $q_N$  expressions give similar results.

An alternate quantification of nonphotochemical fluorescence quenching is as follows (67):

$$NPQ = (F_M - F_M') / F_M'$$

This is a Stern-Volmer formulation that measures the ratio of quenched to remaining fluorescence. The absolute value of NPQ does suffer from distortion by the underlying phycobiliprotein fluorescence, which contributes to both  $F_M$  and  $F_M'$ . This potential problem may be addressed by fluorescence measurements with alternate modulated light sources, such as a blue LED (65), which allow excitation of chlorophyll fluorescence without interference from phycobiliprotein emissions. The use of NPQ also avoids the problematic measurement of  $F_O'$  and is therefore possibly a preferable measure of nonphotochemical quenching in cyanobacteria. Plots of  $q_N$  and NPQ derived from the same fluorescence traces generally show similar although not identical patterns.

Figure 7 presents typical light response curves of nonphotochemical quenching in the cyanobacterium *Synechococcus* sp. strain PCC 7942, along with a curve from a rye plant for comparison. In the cyanobacterium,  $q_N$  (or NPQ) is high in the dark and drops to a minimum near the growth light intensity. In the plant,  $q_N$  climbs steadily as the light intensity surpasses the growth level. These differing patterns reflect a fundamental difference in the predominant processes contributing to nonphotochemical quenching. In plants nonphotochemical quenching is dominated by a mechanism(s) for excitation-dependent thermal dissipation of energy from PS II and its antennae, in competition with fluorescence and photochemistry (1, 46, 47, 57, 66, 67, 153).

In contrast, nonphotochemical quenching in cyanobacteria

largely reflects changes in the PS II fluorescence yield as a result of the state transition mechanism (7, 14, 17, 18, 21, 22, 33, 97, 111, 112), which regulates the distribution of excitation energy between PS II and PS I. The biophysical basis of the energy redistributions remains incompletely understood (2, 17, 18, 118, 156), but it is clear that the relative distribution of excitation energy from the phycobilisome to the two photosystems changes (7, 42, 136). An allophycocyanin-B protein in the phycobilisome core serves as a regulated secondary terminal emitter, which receives about 25% of the excitation energy captured by the phycobilisome. Under state I, this portion of the captured excitation is directed largely to PS II, but under state II, most of it is redirected to PS I, thereby lowering the yield of PS II fluorescence and photochemistry (48, 49, 112, 156). A recent model proposes that reversible changes in the oligomerization of PS II and PS I underpins the state transition mechanism (7). Furthermore, the phycobilisomes diffuse along the surface of the thylakoids (96) sufficiently rapidly that movement of phycobilisomes could be involved in the state transitions.

State transitions in cyanobacteria are regulated by the redox status of the electron transport chain joining PS II and PS I (34, 41, 94, 95, 97, 128, 149). If the chain is reduced, cells tend to state II, with a low yield of PS II fluorescence and a distribution of excitation energy to PS I, which extracts electrons from the chain. If the chain becomes more oxidized the cells shift toward state I, with a higher yield of PS II fluorescence and more distribution of excitation energy to PS II. Although the evidence for redox control of state transitions is quite strong, it is possible to observe conditions of low  $q_P$ , indicating PS II reduction, while cells maintain low  $q_N$ , presumed to reflect state I and an oxidized intersystem transport chain (86a). To accommodate these observations within the framework of redox regulation of state transitions, there must be partial decoupling of PS II and the intersystem redox status under some conditions, or some as yet unknown subtlety to the redox sensing mechanism(s).

Respiratory and photosynthetic electron flow occur via the same electron transport intermediates in cyanobacteria (5, 44, 55, 85, 86, 131, 135, 147) (Fig. 1). Respiratory electron flow in the dark generally poises the electron transport chain toward a reduced state; therefore, in the dark or under very low light, cyanobacteria are in state II (41, 94, 95). This is reflected in low variable fluorescence and high nonphotochemical quenching, as shown in Fig. 7. As light is applied, PS I activity partially oxidizes the electron transport chain and the cells shift toward state I, with higher PS II fluorescence yield (Fig. 2) and lower nonphotochemical quenching (Fig. 7). As light exceeds the growth level, the PS II variable fluorescence yield remains high or drops somewhat (Fig. 2) and nonphotochemical quenching may increase, although not to the levels achieved in darkness (Fig. 7). Finally, if DCMU is added (Fig. 2), PS II centers close, resulting in a rapid rise in fluorescence as photochemical quenching is lost. This rapid rise is equivalent to the fluorescence peak under a brief saturating light pulse, which also closes reaction centers (Fig. 2). The DCMU inhibition of PS II substantially lowers electron input to the transport chain, and the electron transport chain becomes oxidized by continuing PS I activity. This results in a second, slower fluorescence rise phase (Fig. 2) as the cells enter a full state I with maximum PS II fluorescence ( $F_M$ ) (Fig. 2) and minimal nonphotochemical quenching. This redox dependence of the state transition is illustrated in Fig. 8. Titration with the inhibitor DCMU (143) under the growth light intensity progressively blocks PS II activity (Fig. 6), resulting in net oxidation of the transport chain and state I with low  $q_N$  (Fig. 8A). Titration with DBMIB




FIG. 8. Nonphotochemical quenching of fluorescence under DCMU (A) and DBMIB (B) treatments. Modulated fluorescence traces (Fig. 2) were measured under the growth light intensity of  $50 \mu\text{mol of photons m}^{-2} \text{s}^{-1}$ . Results are means and standard errors of measurements on the same culture at  $37^\circ\text{C}$  ( $n = 3$ ). Modified from reference 22 with permission of the publisher.

(143) also inhibits electron transport, but PS II itself remains active, resulting in reduction of the portion of the transport chain preceding the DBMIB binding site on the cytochrome *b*<sub>f</sub> complex. The state transition mechanism senses this change as an apparent overexcitation of PS II, and the cells enter state II with high  $q_N$  (Fig. 8B).

The origin of cyanobacterial nonphotochemical quenching in the state transition mechanism is illustrated in Fig. 9, which compares the 77K fluorescence emission spectra and  $q_N$  values of a cyanobacterial sample in the darkness, under growth light intensity, and after addition of DCMU. The 77K emission spectra, measured with excitation of the phycobilisome at 574 nm, show the changing distribution of energy. In the dark, PS II fluorescence is low, reflecting the high nonphotochemical quenching. Upon illumination,  $q_N$  drops as the excitation energy is redistributed in favour of PS II. This process continues to an extreme upon addition of DCMU.

In plants, excitation-dependent quenching driven by the *trans*-thylakoid  $\Delta\text{pH}$  gradient is the major component of  $q_N$  (1, 46, 47, 57), with only minor contributions from state transitions and other mechanisms (66). In plants, nonphotochemical quenching drops upon collapse of the *trans*-thylakoid pH gradient (67, 153). In cyanobacteria, nonphotochemical quenching does not collapse upon addition of carbonyl cyanide




FIG. 9. 77K Fluorescence emission spectra show that changing  $q_N$  reflects state transitions. The dashed lines emphasize increasing PS II fluorescence and decreasing PS I fluorescence in parallel with the drop in nonphotochemical quenching, measured in the same cultures. A *Synechococcus* sp. strain PCC 7942 culture sample was incubated in the PAM cuvette in darkness, under the growth light intensity ( $50 \mu\text{mol}$  of photons  $\text{m}^{-2} \text{s}^{-1}$ ), and after addition of DCMU under continuing illumination. For each condition,  $q_N$  was measured and a small sample was taken for 77K fluorescence emission spectral analysis. Excitation of 77K fluorescence was carried out with 574 nm light absorbed by the phycobilisome (PBsome). Fluorescence spectra are from samples of equal chlorophyll content but are not otherwise normalized.

*p*-trifluoromethoxyphenylhydrazone (FCCP), an ionophore uncoupler (22, 86a). We therefore concluded that energy-dependent quenching is not a significant contributor to  $q_N$  in cyanobacteria. Interestingly, Delphin et al. (32) present evidence that in red algae, whose chloroplasts also contain phycobilisomes, a plant-type  $\Delta\text{pH}$ -dependent nonphotochemical quenching occurs even under low light levels, in contrast to the state transitions observed in cyanobacteria, which do not depend on  $\Delta\text{pH}$ .

Nonphotochemical quenching reflects changes in PS II photochemistry but not necessarily net energy dissipation from the photosynthetic system, if energy is redirected from PS II to PS I. In cyanobacteria, therefore, a high  $q_N$  does not necessarily mean a low overall photosynthetic efficiency (21, 138). Indeed, redistribution of energy from the PS II-phycobilisome supra-complex to PS I is an important regulatory mechanism in cyanobacteria, to accommodate changing excitation (21, 132, 133) or requirements for ATP to accumulate  $\text{CO}_2$  and nutrients (5, 91, 116). In contrast to this work, nonphotochemical quenching in a *Microcystis* strain has been interpreted as reflecting thermal energy dissipation of excitation. These cyanobacteria had a sustained content of the carotenoid zeaxanthin (60). Furthermore, some cyanobacterial strains such as *Synechocystis* sp. strain PCC 6803 do not display high nonphotochemical quenching after dark adaptation, indicating that they do not enter state II in the dark (51).

The diversity of cyanobacterial pigment and photosynthetic systems requires further characterization of the various origins and regulation of nonphotochemical quenching, particularly under natural growth conditions and limiting nutrient conditions, which can also lead to the loss or alteration of the state transition response (29, 39).

Recent evidence suggests that state II in dark-adapted cyanobacteria might involve an inactivation of the water-splitting complex (83). This limits the electron supply to the  $\text{P}_{680}$  chl of PSII and favors recombination in the reaction center, lowering




FIG. 10. Near the growth light intensity,  $q_N$  reaches a minimum for a wide range of cyanobacterial strains and culture conditions. Mean values are plotted for strains grown at 5 ( $n = 3$ ), 10 ( $n = 5$ ), 15 ( $n = 6$ ), 35 ( $n = 1$ ), 50 ( $n = 20$ ) and 150 ( $n = 3$ )  $\mu\text{mol}$  of photons  $\text{m}^{-2} \text{s}^{-1}$ . The strains were *Anabaena/Nostoc* sp. strain 7120, *Calothrix* sp. strain PCC 7601, *Nostoc* sp., *Pseudanabaena* sp. strain PCC 6901, *Synechococcus* sp. strain PCC 7942, *Synechococcus* sp. strain PCC 6301, and *Synechocystis* sp. strain PCC 6701. Modified from reference 22 with permission of the publisher.

the fluorescence yield (112a). This interpretation suggests that state transitions involve several different mechanisms, since there is clear evidence that under some conditions redirection of excitation energy is responsible for the changes in fluorescence yield observed as state transitions (see, e.g., reference 21).

A methodological problem in measuring light response curves of  $q_N$  is the potential photoinhibitory loss of variable fluorescence as the actinic light is increased above the acclimated growth level. This can be detected if the  $F_M$  level measured with DCMU is lower than a previous  $F_M'$  level, usually that measured around the growth light intensity. In this case, the uninhibited  $F_M$  level for the sample is unknown, and therefore absolute values of  $q_N$  cannot be determined. Nevertheless, if a relative  $q_N$  is calculated by using the highest  $F_M'$  level achieved, the pattern with respect to light intensity is valid. Photoinhibition during measurement can be avoided by minimizing saturating flashes, keeping exposure to high actinic light brief, and terminating light response curves at 5 to 10 times the growth light intensity.

#### Predicting the Acclimated Light Intensity from $q_N$ Light Response

Figure 10 shows that in a wide range of cyanobacteria with different pigment contents and morphologies, grown under different conditions and light histories,  $q_N$  reaches a minimum near the acclimated growth light intensity (22). Therefore, the light intensity to which the population is photosynthetically acclimated can be predicted from a readily measured light response curve of  $q_N$  (or NPQ). Although ambient light is readily measured under many circumstances, this relation between  $q_N$  and growth light shows the range of the overall light regime which is exploited for acclimated growth. In many cases, the natural light regime is highly variable and cyanobacteria must integrate light information over time to regulate synthesis of the abundant proteins of the photosynthetic system. We hope that the  $q_N$  light response curve will prove useful with samples where the past light regime is unknown or samples from variable light regimes in which the optimal light

intensities for acclimated growth are unknown (132, 133). For controlled light acclimation studies, tracking the  $q_N$  minimum over time may show the point at which a population completes acclimation after a light shift. Furthermore, for prediction of electron transport in cyanobacteria from fluorescence parameters, it is essential to measure fluorescence under approximately the acclimated growth light intensity (see below).

Note that this relation does not involve the absolute levels of  $q_N$  or NPQ but simply their pattern in response to light intensity. Determination of actual  $q_N$  or NPQ levels requires the measurement of  $F_M$  by destructive DCMU treatment. If a nondestructive measurement is required and the absolute levels of the parameters are not critical, a simple plot of  $F_M'$  or  $F_V'/F_M'$  against light intensity will suffice; the light intensity at which maximum values are achieved approximately with the acclimated light intensity.

### Inorganic Carbon Accumulation and Fluorescence Quenching

Cyanobacteria and some algae accumulate an intracellular pool of inorganic carbon to limit photorespiration (4, 109). The size of the pool influences both photochemical and nonphotochemical fluorescence quenching (5, 87–91). These fluorescence effects can be used to monitor the transport and accumulation of inorganic carbon noninvasively and in real time (30, 90). One mechanism for the fluorescence effects appears to be a bicarbonate-dependent stimulation of linear electron flow from PS I to O<sub>2</sub>, CO<sub>2</sub>, or nitrite in some strains, which increases photochemical quenching (5, 51, 77, 78, 88, 91, 92).

Carbon accumulation can also influence nonphotochemical quenching, possibly by driving a transition to state II to increase PS I cyclic electron transport to fulfill the need for ATP (93) to drive the accumulation pump.

Miller et al. (89) also describe a form of nonphotochemical quenching which depends on inorganic carbon accumulation in *Synechococcus* sp. strain PCC 7942 cells grown at high light intensities but which does not occur in cells grown at low light intensities. The modulated fluorescence trace from the cells grown at high light intensities resembles a transition to state II upon carbon accumulation, with a drop in  $F_M'$  fluorescence yield and an increase in  $q_N$ . This increase in  $q_N$  was not, however, reflected in a comparable change in the cellular fluorescence emission spectra at 77K. This is in contrast to a large drop in 77K PS II fluorescence emission upon dark adaptation of the same cells, similar to that presented in Fig. 9. This component of carbon accumulation-dependent nonphotochemical quenching in cells grown at high light intensities thus appears distinct from the state transition mechanism. It might ultimately relate to bicarbonate-dependent changes in the PS II water splitting-complex (112a).

### State Transitions Can Be Measured Nondestructively

The change in fluorescence yield during state transitions is strongly influenced by the excitation light, the rate of respiration, the cellular iron supply (39), and the circadian status of the cell (83). Therefore, determining the size of state transitions can provide information about these factors. The magnitude of state transitions can be measured by comparing  $q_N$  or NPQ values under different conditions, but explicit calculation of these parameters requires a lethal DCMU treatment to determine  $F_M$ . This precludes repeated measures on a single sample over time. Alternately, changes in  $q_N$  between different conditions, for example cells in the dark and under illumination, can be quantified by a nondestructive method that does not require explicit measurement of  $F_M$ :

$$(1 - q_{N_{\text{light}}})/(1 - q_{N_{\text{dark}}}) = (F_M' - F_O')/(F_{M_{\text{dark}}} - F_O')$$

In general, for the strains we have studied, this ratio is maximal for a given sample if measured by using the growth light intensity. During prolonged dark incubation, this ratio declines toward 1, as the  $F_{M_{\text{dark}}}$  level approaches but does not reach  $F_M'$ . Thus, during dark incubation, the state transition gradually disappears. This probably reflects the progressive consumption of reserves used to support electron transport in the dark (34, 94, 123). The presence or size of state transitions can also be influenced by iron or nitrogen stress (29, 39), opening the possibility of using the state transition as a noninvasive monitor of these aspects of physiology.

### PREDICTING PHOTOSYNTHESIS FROM FLUORESCENCE

For field measurements, a fluorescence-based estimate of electron transport and carbon dioxide fixation is very valuable (104, 134). Fluorescence measurements are possible with dilute samples; unlike gas exchange, they are specific to photobionts and so do not detect interference from heterotrophic respiratory activity in mixed samples or lichens. The fluorescence transients arise largely from PS II, and so calculations based on fluorescence reflect PS II activity and electron transport through PS II. In extrapolating from fluorescence signals to photosynthesis, we therefore rely on a congruence between PS II activity, net electron transport, and overall photosynthesis. Cyanobacteria and cyanolichens have carbon-concentrating mechanisms which suppress the oxygenase reaction of the ribulose-1,5-bisphosphate oxygenase-carboxylase enzyme (4, 5, 87, 90, 91, 108, 110). This simplifies the empirical relation between PS II activity, reflected in fluorescence signals, and gas exchange, as is also the case in C<sub>4</sub> plants (36).

Sundberg et al. (138) have simultaneously measured fluorescence quenching parameters and CO<sub>2</sub> exchange in cyanolichens to develop a model to predict gross photosynthesis from fluorescence parameters. They found the empirical relation

$$P = \phi \text{PS II} \times I_i \times 1 \text{ CO}_2 \text{ fixed/10 photons}$$

where  $P$  = micromoles of CO<sub>2</sub> fixed per square meter per hour,  $\phi \text{PS II} = (F_M' - F_S)/F_M' = (F_V'/F_M')q_P$  (a fluorescence estimate of the photochemical yield of PS II),  $I_i$  = number of incident photons per square meter per hour, and 1 CO<sub>2</sub> fixed/10 photons is an empirical conversion factor.

This predictor gave good estimates of actual CO<sub>2</sub> fixation near the acclimated growth light intensity. Under higher light, the predictor, which is based on light-driven electron flow through PS II, progressively overestimated actual CO<sub>2</sub> fixation. The overestimation probably reflects electron flow back to O<sub>2</sub> under excess light (22a, 88, 131, 147), which maintains PS II centers open but does not contribute to CO<sub>2</sub> fixation. Other workers have found that this flexible electron transport to O<sub>2</sub> interferes with estimates of CO<sub>2</sub> fixation from fluorescence measures (74). In our experiments, we could estimate the acclimated growth light intensity from the light response curve of nonphotochemical quenching. With this light intensity, we made reasonable predictions of the acclimated rate of CO<sub>2</sub> fixation from fluorescence parameters. The general applicability of this approach must, however, be further tested. The 1 CO<sub>2</sub> fixed/10 photons is an empirical conversion factor that reflects the (unknown) quantum yield of CO<sub>2</sub> fixation and also compensates for the downward distortion of  $F_V'/F_M'$  by phycobilisome fluorescence.

We developed a similar empirical relation to predict gross oxygen evolution from  $\phi \text{PS II}$  in liquid cyanobacterial cultures:




FIG. 11.  $\phi PS II$  reflects  $O_2$  evolution. *Synechococcus* sp. strain PCC 7942 was grown under  $50 \mu\text{mol}$  of photons  $\text{m}^{-2} \text{s}^{-1}$  at  $37^\circ\text{C}$  and then incubated under a range of light intensities. Gross oxygen evolution (micromoles of  $O_2$  per milligram of chlorophyll per hour) was estimated as light-dependent oxygen evolution minus dark uptake.  $\phi PS II = (F_M' - F_S)/F_M' = (F_V'/F_M')q_P$ .  $I_i$  = micromoles of photons incident on the sample per hour.

$$\text{Oxygen evolution} = \phi PS II \\ \times I_i \times 1 O_2/12 \text{ photons} \times 1/\text{chl}$$

where oxygen evolution is expressed as micromoles of  $O_2$  per milligram of chlorophyll per hour,  $\phi PS II = (F_M' - F_S)/F_M' = (F_V'/F_M')q_P$ ,  $I_i$  = number of micromoles of photons incident

per hour;  $1 O_2/12$  photons is an empirical conversion factor, and chl is the chlorophyll content in milligrams.

Figure 11 shows that this relation gives a good approximation of measured oxygen evolution at or near the growth light intensity. At higher light intensities, the predictor increasingly overestimates measured oxygen evolution, again because of pseudocyclic electron flow, with electrons extracted from water by PS II ultimately reaching oxygen as a terminal acceptor. The empirical conversion factor of  $1 O_2/12$  photons again combines the unknown quantum yield of  $O_2$  evolution and compensation for the low  $F_V'/F_M'$  values in cyanobacteria.

In summary, a two-step process gives reasonable estimates of oxygen evolution or carbon fixation under the acclimated growth light intensity. First, a light response curve of nonphotochemical quenching shows the acclimated light intensity. Then oxygen evolution or carbon fixation is estimated from the fluorescence parameter  $\phi PS II$ , the acclimated light level, and an empirical conversion factor calibrated against gas exchange measurements. Failure to measure near the growth light intensity, or an inappropriate empirical calibration factor, can lead to a significant overestimation of actual photosynthesis.

#### APPLYING FLUORESCENCE ANALYSIS TO DIFFERENT CYANOBACTERIA

We have tested the methods described in nine strains of cyanobacteria representing a wide range of pigment contents, phycobilisome structures, and physiological properties, as well as six strains of cyanobacterial lichens, as outlined in Table 2. The methods have been extensively validated with *Synechococcus* sp. strain PCC 7942 (*Anacystis nidulans* R2) grown under a range of light intensities,  $CO_2$  levels, and temperatures (22–24, 26–28; see above). Furthermore, a mutant of this strain lacking phycobilisome rods and containing no PC proved amenable to quenching analysis (Fig. 3) (158). The closely related strain *Synechococcus* sp. strain PCC 6301 (*Anacystis nidulans*) dis-

TABLE 2. Cyanobacteria and cyanolichens used for the modulated fluorescence analyses described in this review

Species	Growth light	Phycobiliprotein <sup>a</sup>
<b>Cyanobacteria</b>		
<i>Anabaena/Nostoc</i> sp. strain PCC 7120	Fluorescent ( $50 \mu\text{mol}$ of photons $\text{m}^{-2} \text{s}^{-1}$ )	APC, PC, PEC
<i>Calothrix/Tolythrix/Fremyella</i> sp. strain PCC 7601	Fluorescent ( $10 \mu\text{mol}$ of photons $\text{m}^{-2} \text{s}^{-1}$ )	APC, PC, PE
<i>Calothrix/Tolythrix/Fremyella</i> sp. strain PCC 7601	Green ( $15 \mu\text{mol}$ of photons $\text{m}^{-2} \text{s}^{-1}$ )	APC, PC, PE
<i>Calothrix/Tolythrix/Fremyella</i> sp. strain PCC 7601	Red ( $15$ and $35 \mu\text{mol}$ of photons $\text{m}^{-2} \text{s}^{-1}$ )	APC, PC
<i>Gloeobacter violaceus</i> PCC 7421	Fluorescent ( $10 \mu\text{mol}$ of photons $\text{m}^{-2} \text{s}^{-1}$ )	APC, PC, PE, PUB
<i>Pseudanabaena</i> sp. strain PCC 6901	Fluorescent ( $50 \mu\text{mol}$ of photons $\text{m}^{-2} \text{s}^{-1}$ )	APC, PC
<i>Synechococcus</i> sp. strain PCC 7942 ( <i>Anacystis nidulans</i> R2)	Incandescent ( $10, 15, 50,$ and $150 \mu\text{mol}$ of photons $\text{m}^{-2} \text{s}^{-1}$ )	APC, PC
<i>Synechococcus</i> sp. strain PCC 7942 phycobilisome-minus	Incandescent ( $50 \mu\text{mol}$ of photons $\text{m}^{-2} \text{s}^{-1}$ )	APC
<i>Synechococcus</i> sp. strain PCC 6301 ( <i>Anacystis nidulans</i> )	Fluorescent ( $10$ and $50 \mu\text{mol}$ of photons $\text{m}^{-2} \text{s}^{-1}$ )	APC, PC
<i>Synechocystis</i> sp. strain PCC 6701	Fluorescent ( $50 \mu\text{mol}$ of photons $\text{m}^{-2} \text{s}^{-1}$ )	APC, PC, PE
<i>Nostoc</i> from lichen <i>Peltigera canina</i>	Fluorescent ( $5 \mu\text{mol}$ of photons $\text{m}^{-2} \text{s}^{-1}$ )	APC, PC, (PE, PEC?)
<b>Cyanolichens (Nostoc)</b>		
<i>Leptogium coralloideum</i>	Natural	APC, PC, (PE, PEC?)
<i>Lobaria scrobiculata</i>	Natural	APC, PC, (PE, PEC?)
<i>Nephroma bellum</i>	Natural	APC, PC, (PE, PEC?)
<i>Peltigera malacea</i>	Natural	APC, PC, (PE, PEC?)
<i>Peltigera neopolydactyla</i>	Natural	APC, PC, (PE, PEC?)
<i>Nephroma arcticum</i> ( <i>Nostoc</i> and green algae <i>Coccomyxa</i> )	Natural	APC, PC, (PE, PEC?)

<sup>a</sup> APC, allophycocyanin; PC, phycocyanin; PE, phycoerythrin; PEC, phycoerythrocyanin; PUB, phycoerythrin containing phycocourobilin chromophores.

plays very similar fluorescence properties (21a). Both these strains have a somewhat unusual phycobilisome with a core composed of only two cylinders.

*Calothrix* sp. strain PCC 7601 is a heterocystous, filamentous strain which strongly regulates its content of phycoerythrin and phycocyanin according to the relative supply of green and red light. Quenching analysis has shown that when this strain is grown under green light, energy captured by phycoerythrin is transferred from the phycobilisome to PS I to maintain balanced electron transport. This transfer is reflected in a high  $q_N$  and serves as a good demonstration that in a cyanobacterium a high  $q_N$  does not usually reflect excitation dissipation but, rather, reflects the transfer of excitation to PS I at the expense of PS II. Upon transfer to red light or prolonged growth under red light, this excitation transfer stops, the PS II fluorescence yield increases, and  $q_N$  drops (21).

Four other diverse cyanobacterial strains have also shown responses similar to that of *Synechococcus* sp. strain PCC 7942 (21a): *Anabaena/Nostoc* sp. strain PCC 7120 is a heterocystous, filamentous strain with another atypical phycobilisome structure (35), *Nostoc* sp. is a strain originally isolated from the lichen *Peltigera canina*, *Synechocystis* sp. strain PCC 6701 is a unicellular strain which contains phycoerythrin; and *Pseudanabaena* sp. strain PCC 6901 is a gas-vacuolated strain forming short filaments.

Six lichens with *Nostoc* strains as symbionts have also proved amenable to quenching analysis by using a Hansatech cuvette designed for leaf discs; *Leptogium coralloideum*, *Lobaria scrobiculata*, *Nephroma bellum*, *Peltigera malacea*, *Peltigera neopolydactyla*, and the cephalodia regions of *Nephroma arcticum*, a tri-partite lichen with the green algae *Coccomyxa* and *Nostoc*. In lichens, the dark-light state II-state I transition tends to be small, probably reflecting fungus-cyanobacterium interactions in respiration and carbohydrate consumption (138).

Goosney and Miller (51) found that the widely studied facultative heterotrophic strain *Synechocystis* sp. strain PCC 6803 shows little or no increase in  $F_M'$  upon illumination under most conditions. The fluorescence induction trace in this strain is rather plant-like in that  $F_M'$  measured in dark-adapted cells is close to  $F_M$ . This distinct pattern might reflect differences in this strain in the redox balance of intersystem electron transport in the dark, such that the cells do not enter state II in the dark. This plant-like induction pattern upon illumination can occur in other strains, particularly when dark respiration is slow as under conditions of low carbohydrate reserves or under nutrient stress (29, 39). Alternatively, there may be more fundamental distinctions in the organization of light capture and electron transport between various strains.

Interestingly, the unusual cyanobacteria *Gloeobacter violaceus* PCC 7421 was the only strain surveyed for which quenching analysis proved impossible. This strain lacks thylakoids (115) and instead of typical phycobilisomes contains simpler rod-like phycobiliprotein structures (52) associated with the plasma membrane. In our measurements, this strain showed almost no variable fluorescence even when growth and oxygen evolution were readily measurable (21a). 77K fluorescence spectra from this strain also lack the expected long-wavelength PS I emission, even though the presence of PS I was verified functionally (64). Clearly, the unusual photosynthetic system of this cyanobacterium results in distinct fluorescence properties.

## CONCLUSIONS AND PROSPECTS

### Tracking Acclimation Status in the Laboratory and Field

Fluorescence analysis is an integral part of studies of photosynthesis in cyanobacteria and other organisms. In recent years, advances in instrumentation and interpretation have greatly expanded the applications of fluorescence to ecophysiological and molecular studies (6, 13, 40, 99).

For cyanobacteria, estimating the acclimated light in a population from the light response of nonphotochemical quenching (22) will allow rapid tracking of acclimation in laboratory experiments or field studies. Cells must integrate changing environmental signals (132, 133) to regulate the expression of abundant proteins such as ribulose-1,5-bisphosphate carboxylase/oxygenase or phycobilisomes, in order to produce appropriate long-term levels of protein. The use of fluorescence to monitor acclimation (10, 61, 73, 74, 130) may show how cyanobacteria and cyanolichens set their targets for gene expression and metabolic acclimation in the face of changing light, environmental factors, and circadian status.

The size of the dark-to-light state transition is strongly influenced by cellular respiration, which poises the electron transport chain (34, 55, 84–86, 94, 95, 131, 135). Therefore, the state transition may provide estimates of respiration and indirectly of the reserves available to support respiration. Further work in this area might lead to rapid measures of the levels of reserves in cyanobacteria or cyanolichens, as they fluctuate diurnally or seasonally (123, 124). Estimates of reserves are valuable in understanding physiological and differentiation responses of cyanobacteria and also in determining production available for export from the community.

The size of the dark-to-light state transition is also strongly influenced by cellular iron and nitrogen status (29, 39). Under iron limitation, which is widespread in nature (8), cyanobacteria produce alternate chlorophyll-protein complexes associated with PS II (70, 75, 76), which leads to suppression of the state transition (39). Therefore, the size of state transition may be useful as a measure of iron limitation.

The iron example illustrates that as we apply fluorescence analysis to natural cyanobacteria, nutrient or reserve limitations may greatly alter fluorescence signals from those observed with laboratory cultures. Thus, further work is needed on measuring and interpreting fluorescence signals under non-optimal and natural conditions, where pigment composition and cellular organization may be very different from typical laboratory cultures. Simultaneous parallel detection of fluorescence transients at several excitation and emission wavelengths (61, 65) holds great promise for the study of mixed populations or strains with complex pigment compositions.

### Conclusions

Chlorophyll fluorescence signals from cyanobacteria and cyanolichens show patterns very distinct from those of green plants. Therefore, the fluorescence measurements and analyses originally developed for green plants must be modified, but cyanobacterial fluorescence also yields information not accessible from plant fluorescence signals. The cellular phycobiliprotein content influences the  $F_O$  level fluorescence, particularly when phycobiliprotein levels are high. This leads in some cases to downward distortion of the parameter  $F_V/F_M$ , which is widely used as an index of PS II activity. The photochemical quenching coefficient,  $q_P$ , provides a robust index of the balance between excitation of PS II and electron transport and shows that cyanobacteria have a high and flexible capacity to remove electrons from PS II. Nonphotochemical quenching of

PS II variable fluorescence in cyanobacteria reflects the state transition mechanism for distribution of excitation between the photosystems. A characteristic decline of nonphotochemical quenching during a shift from dark to increasing light provides a valuable means of estimating the light level to which a cyanobacterial population is photosynthetically acclimated. Gross oxygen evolution or carbon fixation can be estimated in at least some cases (138; but see reference 74)) from the fluorescence parameter  $\Phi_{PS II}$ , the acclimated growth light level, and an empirically verified apparent quantum yield.

## APPENDIX

### Cyanobacterial Cultures and Pigment Measurements

Cyanobacteria were grown in BG-11 inorganic medium (114), supplementally buffered with 10 mM 3-(*N*-morpholino)propanesulfonic acid (MOPS) to a final pH of 7.5. *Synechococcus* sp. strain PCC 7942 cultures (300 ml) were grown in flat flasks bubbled with 5% CO<sub>2</sub> in air (about 1 ml s<sup>-1</sup>) at 37 or 25°C with continuous, even illumination of 10, 15, 50, or 150 μmol of photons m<sup>-2</sup> s<sup>-1</sup>. For *Synechococcus* sp. strain PCC 7942, the chlorophyll and phycocyanin contents were determined by using whole-cell spectra as described by Myers et al. (98), corrected for scattering by subtracting the absorbance at 750 nm from the chlorophyll and phycocyanin peaks. Whole-cell spectra are less useful for filamentous strains, since significant and variable light scattering distorts spectra. For these strains, chlorophyll was extracted in methanol and measured by the method of Tandeau de Marsac and Houmard (140). For the other strains used in the  $q_N$  measurements shown in Fig. 9, the growth conditions were generally similar but the temperature ranged from 18 to 37°C, the light intensity ranged from 5 to 150 μmol of photons m<sup>-2</sup> s<sup>-1</sup>, either fluorescent or incandescent, and the CO<sub>2</sub> supply ranged from ambient to 5% in air.

### Modulated Fluorometer Configuration and Measurement Procedure

In our experiments, chlorophyll *a* fluorescence induction was measured with a pulse amplitude modulated fluorometer (128, 129) (PAM chlorophyll fluorometer; Walz, Effeltrich, Germany) with the PAM 103 accessory and a Schott KL1500 lamp (Schott, Mainz, Germany) to provide saturating flashes. The recording device, in our case a chart recorder, should respond to changes within less than 100 ms (128). A PAM-compatible system of cuvette, magnetic stirrer, oxygen electrode, and halogen incandescent actinic lamp were used for the simultaneous measurement of fluorescence and oxygen evolution (Hansatech, King's Lynn, United Kingdom) (151). Simultaneous measurement strengthens the conclusions which can be drawn, since the two techniques are complementary. This cuvette system allows measurement of liquid samples down to concentrations of about 0.2 μg of chlorophyll/ml, suitable for laboratory cultures or somewhat concentrated natural samples. An alternate cuvette (Walz ED-101 US) (125) uses a smaller vessel and light guides to extend the functional concentration range to less than 1 μg of chlorophyll per liter, allowing direct measurement of many natural water samples. The actinic beam and fluorescence detector are set at 90° in this system, rather than at 180° as in the Hansatech cuvette. This alternate geometry may result in changes in some measurements (60a), and so the influences of cuvette geometry on signals are an issue for further research. A further innovation is a system involving light-emitting diodes of different wavelengths, for preferential

excitation of specific pigments (65). This technique shows promise for resolution of fluorescence signals from different photobionts in mixed samples. Comparable modulated fluorometers, with variations in the nature and wavelength of the measuring and actinic lights and in the cuvettes, are available from several companies.

Culture samples, generally at around 2 μg of chlorophyll per ml, were dark adapted for 5 min in the Hansatech cuvette. The analysis procedure is outlined in Fig. 2. Minimum fluorescence,  $F_O$ , was determined by illuminating the dark-adapted cells with a low-intensity light modulated at 1.6 kHz (average intensity, 0.14 μmol of photons m<sup>-2</sup> s<sup>-1</sup>) from a light-emitting diode (peak emission, 655 nm). Fluorescence was detected at wavelengths greater than 700 nm. The intensity of the measuring beam should be checked to ensure that it is sufficiently weak to avoid electron transport, detected as changes in the oxygen electrode response (20). A 1-s flash of saturating white light (8,000 μmol of photons m<sup>-2</sup> s<sup>-1</sup>) was then given to determine the maximal fluorescence in the dark-adapted state,  $F_{M\text{dark}}$ , with all PS II centers closed by the saturating flash. In cyanobacteria,  $F_{M\text{dark}}$  is generally significantly lower than the maximal fluorescence,  $F_M$  (20, 22, 23, 27, 28, 91), although this is not the case in all strains under all conditions (see, e.g., reference 51). After a further 30 s, the actinic light was activated. Steady-state fluorescence,  $F_S$ , was reached within 2 min. Minimum fluorescence in the light-adapted state,  $F_{O'}$ , was then measured by briefly interrupting the actinic beam and either leaving culture in darkness for ca. 5 s or applying weak far-red light (ca. 5 μmol of photons m<sup>-2</sup> s<sup>-1</sup>; >700 nm) to excite PS I activity and extract electrons from the transport chain. Both procedures for measuring  $F_{O'}$  gave similar results in cyanobacterial measurements. The actinic light was then resumed, and after  $F_S$  was reestablished, a saturating light pulse was given to again close all PS II centers, driving photochemical quenching to zero for determination of maximal fluorescence in the light-adapted state,  $F_{M'}$ . The actinic light intensity was then increased, and the process was repeated sequentially to generate a light response curve.

Cyanobacterial cells have CO<sub>2</sub>-concentrating mechanisms (CCM) (4, 87), and when this system is induced the cells do not usually become CO<sub>2</sub> limited during short measurements. Laboratory cyanobacterial cultures are often supplemented with high CO<sub>2</sub> levels, and under these growth conditions the CCM activity is repressed. In these cases, photosynthesis may deplete the dissolved inorganic carbon supply of the cuvette volume, resulting in CO<sub>2</sub> limitation of photosynthesis. Even in cells grown under low inorganic carbon, prolonged photosynthesis in a closed cuvette can deplete the inorganic carbon sufficiently to cause carbon limitation. Furthermore, electron flow to oxygen is often dependent on the presence of an intracellular bicarbonate pool, so CO<sub>2</sub> limitation can also limit electron flow to oxygen as a terminal acceptor (88). The onset of CO<sub>2</sub> limitation results in steady-state  $F_S$  fluorescence increasing toward  $F_{M'}$ , in parallel with a decline in the oxygen evolution rate. Cultures grown under CO<sub>2</sub> supplementation were therefore routinely supplemented with 7 mM NaH<sub>2</sub>CO<sub>3</sub> at the start of the measurement.

Finally, DCMU (final concentration, 0.5 μM) was injected into the cuvette to bind to PS II centers (143). This leads to a rapid rise in fluorescence to a level similar to  $F_{M'}$ , as PS II closes, photochemistry is blocked, and photochemical quenching is lost (Fig. 2). The loss of PS II activity leads to oxidation of the plastoquinone pool, which in turn drives a slower fluorescence rise phase (Fig. 2) as the cells go to full state I with maximal fluorescence,  $F_M$ . The concentration of DCMU required for rapid PS II closure is somewhat dependent on the

strain and growth conditions and should be verified empirically. The parameters  $F_O$ ,  $F_{M\text{dark}}$ ,  $F_S$ ,  $F_O'$ ,  $F_M'$ , and  $F_M$  were used for the calculation of photochemical ( $q_P$ ) and nonphotochemical ( $q_N$ ) quenching (145), and the apparent efficiency of excitation energy capture by open PS II reaction centers ( $F_V'/F_M'$ ) (45).

Figure 2 is a fluorescence trace from our usual protocol for measuring room temperature fluorescence in *Synechococcus* sp. strain PCC 7942, showing a typical cyanobacterial response over a series of increasing light intensities. The number of saturating flashes is kept to a minimum to avoid photoinhibition or sustained induction of electron transport. Schreiber et al. (128) described the use of much shorter flashes (50 ms) which can be applied to measure  $F_M'$  more frequently with a reduced risk of photoinhibition or perturbation of electron transport.

### 77K Fluorescence Emission Spectra

A clear acrylic rod sample holder for 77K fluorescence was dipped briefly into the Hansatech PAM cuvette to collect about 100  $\mu\text{l}$ ; this subsample was plunged directly into liquid nitrogen. The sample holder was attached to a fiberoptic fluorometer, and 77K fluorescence emission spectra were collected (100), with excitation at 574 nm. The 77K spectra are therefore from cells under the same conditions as those used for room temperature fluorescence quenching and oxygen measurements.

### Alternate Fluorescence Quenching Parameters

Havaux et al. (53) and Walters and Horton (152) have proposed the parameter  $(1/F_O) - (1/F_M)$  as a measure of the rate constant of excitation trapping by the PS II reaction center and  $1/F_M$  as a measure of nonphotochemical loss of excitation energy from the light-harvesting antennae. In our experiments, plots of  $(1/F_O) - (1/F_M)$  closely parallel  $F_V'/F_M$ . The  $1/F_M$  parameter shows some discrepancy from cyanobacterial  $q_N$  and NPQ, possibly reflecting the distinct nature of nonphotochemical quenching in cyanobacteria and plants.

An alternate approach to extracting information from fluorescence signals is the pump and probe method reviewed by Falkowski and Kolber (40). For methodological and theoretical reasons, these authors define an alternate quantum efficiency term,

$$\Delta\phi_{\text{sat}} = (F_{\text{sat}} - F_{\text{probe}})/F_{\text{probe}}$$

as the difference in fluorescence yield driven by a weak probe light before and immediately following a saturating flash, divided by the fluorescence yield before the flash. This alternate parameter can be related to  $F_V'/F_M$  as

$$F_V'/F_M = \Delta\phi_{\text{sat}}/(\Delta\phi_{\text{sat}} + 1)$$

Another approach developed particularly for oceanographic purposes is fast repetition rate fluorescence (40), which uses a rapid train of 20 to 60 brief (5- $\mu\text{s}$ ), nonsaturating pulses applied at frequencies up to 200 kHz. Fluorescence excited by the first flash gives the  $F_O$  level fluorescence with all reaction centers open. The rapid succession of flashes then progressively closes PS II centers, driving fluorescence toward  $F_M$ . The rate of saturation is proportional to the effective absorption cross-section of PS II,  $\sigma_{\text{PS II}}$ , a useful parameter which is not determined by methods using single saturating pulses. The  $F_M$  measured with this train of pulses is not equivalent to the  $F_M$  determined with a single saturating pulse used in the modulated fluorometer (127, 128) or the pump and probe methods.

Therefore, data from fast repetition rate fluorescence are not directly comparable or convertible to data from the other methods. Nevertheless, the results generate conclusions which are qualitatively similar to those obtained from quenching analysis involving modulated fluorescence (127).

### ACKNOWLEDGMENTS

We thank our collaborators Doug Bruce, Amanda Cockshutt, Kristin Palmqvist, Bodil Sundberg, and Guoqing Zhou for valuable discussions. Anthony Miller, Darwyn Coxson, Henrik Schubert, Nicole Tandeau de Marsac, and Jean Houmar have also contributed generously to discussions of this work.

This work was supported by the Swedish Natural Science Research Council grants to G.Ö.

### REFERENCES

1. Adams, W. W., and B. Demmig-Adams. 1993. Energy dissipation and photoprotection in leaves of higher plants, p. 27–36. In H. Y. Yamamoto and C. M. Smith (ed.), Photosynthetic responses to the environment. American Society of Plant Physiologists, Rockville, Md.
2. Allen, J. F., C. E. Sanders, and N. G. Holmes. 1985. Correlation of membrane protein phosphorylation with excitation energy distribution in the cyanobacterium *Synechococcus* 6301. FEBS Lett. **193**:271–275.
3. Asada, K., K. Kiso, and K. Yoshikawa. 1974. Univalent reduction of molecular oxygen by spinach chloroplasts on illumination. J. Biol. Chem. **249**:2175–2181.
4. Badger, M. R., and G. D. Price. 1992. The CO<sub>2</sub> concentrating mechanism in cyanobacteria and green algae. Physiol. Plant. **84**:606–615.
5. Badger, M. R., and U. Schreiber. 1993. Effects of inorganic carbon accumulation on photosynthetic oxygen reduction and cyclic electron flow in the cyanobacterium *Synechococcus* PCC 7942. Photosynth. Res. **37**:177–191.
6. Baker, N. R., P. K. Farage, C. M. Stirling, and S. P. Long. 1994. Photoinhibition of crop photosynthesis in the field at low temperatures, p. 349–363. In N. R. Baker and J. R. Bowyer (ed.), Photoinhibition of photosynthesis: from molecular mechanisms to the field. Bios Scientific Publishers, Oxford, United Kingdom.
7. Bald, D., J. Kruij, and M. Rögner. 1996. Supramolecular architecture of cyanobacterial thylakoid membranes: how is the phycobilisome connected with the photosystems? Photosynth. Res. **49**:103–118.
8. Behrenfeld, M. J., A. J. Bale, Z. S. Kolber, J. Aiken, and P. G. Falkowski. 1996. Confirmation of iron limitation of phytoplankton photosynthesis in the equatorial pacific-ocean. Nature **383**:508–511.
9. Bilger, W., and U. Schreiber. 1986. Energy-dependent quenching of dark-level chlorophyll fluorescence in intact leaves. Photosynth. Res. **10**:303–308.
10. Bilger, W., B. Budel, R. Mollenhauer, and D. Mollenhauer. 1994. Photosynthetic activity of 2 developmental stages of a *Nostoc* strain (cyanobacteria) isolated from *Geosiphon pyriforme* (mycota). J. Phycol. **30**:225–230.
11. Binder, A., R. Hauser, and D. Krogmann. 1984. Respiration in energy transducing membranes of the thermophilic cyanobacterium *Mastigocladus laminosus*. I. Relation of the respiratory and photosynthetic electron transport. Biochim. Biophys. Acta **765**:241–246.
12. Björkman, O., and B. Demmig. 1987. Photon yield of O<sub>2</sub> evolution and chlorophyll fluorescence characteristics at 77K among vascular plants of diverse origins. Planta **170**:489–504.
13. Bolhär-Nordenkamp, H. R., and G. Öquist. 1993. Chlorophyll fluorescence as a tool in photosynthesis research, p. 193–206. In D. O. Hall, J. M. O. Scurlock, H. R. Bolhär-Nordenkamp, R. C. Leegood, and S. P. Long (ed.), Photosynthesis and production in a changing environment: a field and laboratory manual. Chapman & Hall, London, United Kingdom.
14. Bonaventura, C., and J. Myers. 1969. Fluorescence and oxygen evolution for *Chlorella pyrenoidosa*. Biochim. Biophys. Acta **189**:366–383.
15. Bovy, A., G. de Vrieze, M. Borrias, and P. Weisbeek. 1992. Transcriptional regulation of the plastocyanin and cytochrome c<sub>553</sub> genes from the cyanobacterium *Anabaena* species PCC 7937. Mol. Microbiol. **6**:1507–1513.
16. Briggs, L. M., V. L. Pecoraro, and L. McIntosh. 1990. Copper-induced expression, cloning, and regulatory studies of the plastocyanin gene from the cyanobacterium *Synechocystis* sp. PCC 6803. Plant Mol. Biol. **15**:633–542.
17. Bruce, D., and J. Biggins. 1985. Mechanism of the light-state transition in photosynthesis. V. 77K linear dichroism of *Anacystis nidulans* in State 1 and State 2. Biochim. Biophys. Acta **810**:295–301.
18. Bruce, D., J. Biggins, T. Steiner, and M. Thewalt. 1985. Mechanism of the light-state transition in photosynthesis. IV. Picosecond fluorescence spectroscopy of *Anacystis nidulans* and *Porphyridium cruentum* in State 1 and State 2. Biochim. Biophys. Acta **810**:237–246.
19. Bryant, D. A. (ed.). 1994. The molecular biology of cyanobacteria. Kluwer Academic Publishers, Dordrecht, The Netherlands.
20. Büchel, C., and C. Wilhem. 1993. *In vivo* analysis of slow chlorophyll fluo-

- rescence induction kinetics in algae: progress, problems and perspectives. *Photochem. Photobiol.* **58**:137–148.
21. **Campbell, D.** 1996. Complementary chromatic adaptation alters photosynthetic strategies in the cyanobacterium *Calothrix*. *Microbiology* **142**:1255–1263.
  - 21a. **Campbell, D.** Unpublished data.
  22. **Campbell, D., and G. Öquist.** 1996. Predicting light acclimation in cyanobacteria from non-photochemical quenching of PSII fluorescence, which reflects state transitions in these organisms. *Plant Physiol.* **111**:1293–1298.
  - 22a. **Campbell, D., and G. Öquist.** Unpublished data.
  23. **Campbell, D., D. Bruce, C. Carpenter, P. Gustafsson, and G. Öquist.** 1996. Two forms of the photosystem II D1 protein alter energy dissipation and state transitions in the cyanobacterium *Synechococcus* sp. PCC 7942. *Photosynth. Res.* **47**:131–144.
  24. **Campbell, D., G. Zhou, P. Gustafsson, G. Öquist, and A. K. Clarke.** 1995. Electron transport regulates exchange of two forms of Photosystem II D1 protein in the cyanobacterium *Synechococcus*. *EMBO J.* **14**:5457–5466.
  25. **Campbell, D., J. Houmard, and N. Tandeau de Marsac.** 1993. Electron transport regulates cellular differentiation in the filamentous cyanobacterium *Calothrix*. *Plant Cell* **5**:451–463.
  26. **Clarke, A. K., and D. Campbell.** 1996. Inactivation of the *petE* gene for plastocyanin lowers photosynthetic capacity and exacerbates chilling-induced photoinhibition in the cyanobacterium *Synechococcus*. *Plant Physiol.* **112**:1551–1561.
  27. **Clarke, A. K., D. Campbell, P. Gustafsson, and G. Öquist.** 1995. Dynamic responses of photosystem II and phycobilisomes to changing light in the cyanobacterium *Synechococcus* sp. PCC 7942. *Planta* **197**:553–562.
  28. **Clarke, A. K., V. M. Hurry, P. Gustafsson, and G. Öquist.** 1993. Two functionally distinct forms of the PS II reaction-center protein D1 in the cyanobacterium *Synechococcus* sp. PCC 7942. *Proc. Natl. Acad. Sci. USA* **90**:11985–11989.
  29. **Collier, J. L., S. K. Herbert, D. C. Fork, and A. R. Grossman.** 1994. Changes in the cyanobacterial photosynthetic apparatus during acclimation to macronutrient deprivation. *Photosynth. Res.* **42**:173–183.
  30. **Crotty, C. M., P. N. Tyrrell, and G. S. Espie.** 1994. Quenching of chlorophyll-a fluorescence in response to Na<sup>+</sup>-dependent HCO<sub>3</sub><sup>-</sup> transport-mediated accumulation of inorganic carbon in the cyanobacterium *Synechococcus* UTEX-625. *Plant Physiol.* **104**:785–791.
  31. **Dau, H.** 1994. Molecular mechanisms and quantitative models of variable photosystem II fluorescence. *Photochem. Photobiol.* **60**:1–23.
  32. **Delphin, E., J.-C. Duval, A.-L. Etienne, and D. Kirilovsky.** 1996. State transitions or ΔpH-dependent quenching of photosystem II fluorescence in red algae. *Biochemistry* **35**:9435–9445.
  33. **Demmig-Adams, B., W. W. Adams III, F.-C. Czygan, U. Schreiber, and O. L. Lange.** 1990. Differences in the capacity for radiationless energy dissipation in green and blue-green algal lichens associated with differences in carotenoid composition. *Planta* **180**:582–589.
  34. **Dominy, P. J., and W. P. Williams.** 1987. The role of respiratory electron flow in the control of excitation energy distribution in blue-green algae. *Biochim. Biophys. Acta* **892**:264–274.
  35. **Ducret, A., W. Sidler, and H. Zuber.** 1993. Structural studies on the atypical phycobilisome core complex of the cyanobacterium *Anabaena* sp. PCC 7120, p. 37. *In* 1993 Westcoast Cyanobacterial Workshop.
  36. **Edwards, G. E., and N. R. Baker.** 1993. Can CO<sub>2</sub> assimilation in maize leaves be predicted accurately from chlorophyll fluorescence analysis? *Photosynth. Res.* **37**:89–102.
  37. **Elston, T., H. Wang, and G. Oster.** 1998. Energy transduction in ATP synthase. *Nature* **391**:510–513.
  38. **Evans, E. H., and R. G. Brown.** 1994. An appraisal of photosynthetic fluorescence kinetics as a probe of plant function. *J. Photochem. Photobiol. B* **22**:95–104.
  39. **Falk, S., G. Samson, D. Bruce, N. P. A. Huner, and D. E. Laudenbach.** 1995. Functional analysis of the iron-stress induced CP-43'-polypeptide of PS-II in the cyanobacterium *Synechococcus* sp. PCC-7942. *Photosynth. Res.* **45**:51–60.
  40. **Falkowski, P. G., and Z. Kolber.** 1995. Variations in the chlorophyll fluorescence yields in phytoplankton in the world oceans. *Aust. J. Plant Physiol.* **22**:341–355.
  41. **Fork, D. C., and K. Satoh.** 1983. State I-State II transitions in the thermophilic blue-green alga (cyanobacterium) *Synechococcus lividus*. *Photochem. Photobiol.* **37**:421–427.
  42. **Fujita, Y., A. Murakami, K. Aizawa, and K. Ohki.** 1994. Short-term and long-term adaptation of the photosynthetic apparatus: homeostatic properties of thylakoids, p. 677–692. *In* D. A. Bryant (ed.), *The molecular biology of Cyanobacteria*. Kluwer Academic Publishers, Dordrecht, The Netherlands.
  43. **Furbank, R. T., and M. R. Badger.** 1983. Oxygen exchange associated with electron transport and photophosphorylation in spinach thylakoids. *Biochim. Biophys. Acta* **723**:400–409.
  44. **Geerts, D., H. Schubert, G. de Vrieze, M. Borrias, H. C. P. Matthijs, and P. J. Weisbeek.** 1994. Expression of *Anabaena* PCC 7937 plastocyanin in *Synechococcus* PCC 7942 enhances photosynthetic electron transfer and alters the electron distribution between photosystem I and cytochrome-c oxidase. *J. Biol. Chem.* **269**:28068–28075.
  45. **Genty, B., J. M. Briantais, and N. R. Baker.** 1989. The relationship between the quantum yield of photosynthetic electron transport and quenching of chlorophyll fluorescence. *Biochim. Biophys. Acta* **990**:87–92.
  46. **Gilmore, A. M.** 1997. Mechanistic aspects of xanthophyll cycle dependent photoprotection in higher plant chloroplasts and leaves. *Physiol. Plant.* **99**:197–209.
  47. **Gilmore, A. M., and H. Y. Yamamoto.** 1993. Biochemistry of xanthophyll-dependent nonradiative energy dissipation, p. 160–165. *In* H. Y. Yamamoto and C. M. Smith (ed.), *Photosynthetic responses to the environment*. American Society of Plant Physiologists, Rockville, Md.
  48. **Gindt, Y. M., J. H. Zhou, D. A. Bryant, and K. Sauer.** 1994. Spectroscopic studies of phycobilisome subcore preparations lacking key core chromophores—assignment of excited-state energies to the L(CM), beta(18) and alpha (AP-B) chromophores. *Biochim. Biophys. Acta* **1186**:153–162.
  49. **Glazer, A. N., C. Chan, R. C. Williams, S. W. Yeh, and J. H. Clark.** 1985. Kinetics of energy flow in the phycobilisome core. *Science* **230**:1051–1053.
  50. **Goetze, D. C., and R. Carpentier.** 1994. Ferredoxin-NADP<sup>+</sup> reductase is the site of oxygen reduction in pseudocyclic electron transport. *Can. J. Bot.* **72**:256–260.
  51. **Goosney, D. L., and A. G. Miller.** 1997. High rates of O<sub>2</sub> photoreduction by the unicellular cyanobacterium *Synechocystis* PCC 6803 as determined by the quenching of chlorophyll fluorescence. *Can. J. Bot.* **75**:394–401.
  52. **Guglielmi, G., G. Cohen-Bazire, and D. A. Bryant.** 1981. The structure of *Gloeobacter violaceus* and its phycobilisomes. *Arch. Microbiol.* **129**:181–189.
  53. **Havaux, M., R. J. Straser, and H. Greppin.** 1991. A theoretical and experimental analysis of the qP and qN coefficients of chlorophyll fluorescence quenching and their relation to photochemical and nonphotochemical events. *Photosynth. Res.* **27**:41–55.
  54. **Herbert, S. K., R. E. Martin, and D. C. Fork.** 1995. Light adaptation of cyclic electron-transport through photosystem-I in the cyanobacterium *Synechococcus* sp. PCC7942. *Photosynth. Res.* **46**:277–285.
  55. **Hirano, M., K. Satoh, and S. Katoh.** 1980. Plastoquinone as a common link between photosynthesis and respiration in blue-green alga. *Photosynth. Res.* **1**:149–162.
  56. **Holzwarth, A. R.** 1993. Is it time to throw away your apparatus for chlorophyll fluorescence induction? *Biophys. J.* **64**:1280–1281.
  57. **Horton, P., A. V. Ruban, and R. G. Walters.** 1996. Regulation of light harvesting in green plants. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **47**:655–684.
  58. **Hovenden, M. J., and R. D. Seppelt.** 1995. Utility of modulated fluorescence in measuring photosynthetic activity of Antarctic plants: field and laboratory studies. *Aust. J. Plant Physiol.* **22**:321–330.
  59. **Huner, N. P. A., G. Öquist, V. M. Hurry, M. Krol, S. Falk, and M. Griffith.** 1993. Photosynthesis, photoinhibition and low temperature acclimation in cold tolerant plants. *Photosynth. Res.* **37**:19–39.
  60. **Ibelings, B. W., B. M. A. Kroon, and L. R. Mur.** 1994. Acclimation of Photosystem-II in a cyanobacterium and a eukaryotic green-alga to high and fluctuating photosynthetic photon flux densities, simulating light regimes induced by mixing in lakes. *New Phytol.* **128**:407–424.
  - 60a. **Ivanov, A., and N. Huner.** Personal communication.
  61. **Jensen, M., and K. Siebke.** 1997. Fluorescence imaging of lichens in the macro scale. *Symbiosis* **23**:183–195.
  62. **Jones, L. W., and J. Myers.** 1963. A common link between photosynthesis and respiration in a blue-green alga. *Nature* **199**:670–672.
  63. **Kana, T. M.** 1993. Rapid oxygen cycling in *Trichodesmium thiebautii*. *Limnol. Oceanogr.* **38**:18–24.
  64. **Koenig, F., and M. Schmidt.** 1995. *Gloeobacter violaceus*—investigation of an unusual photosynthetic apparatus—absence of the long-wavelength emission of photosystem-I in 77 K fluorescence-spectra. *Physiol. Plant.* **94**:621–628.
  65. **Kolbowski, J., and U. Schreiber.** 1995. Computer controlled phytoplankton analyser based on chlorophyll fluorescence analysis using 4 different wavelengths. *Xth International Photosynthesis Congress. Photosynth. Res.* **51**:10–013.
  66. **Krause, G. H., and E. Weis.** 1991. Chlorophyll fluorescence and photosynthesis: the basics. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **42**:313–349.
  67. **Krause, G. H., C. Vernotte, and J.-M. Briantais.** 1982. Photoinduced quenching of chlorophyll fluorescence in intact chloroplasts and algae. Resolution into two components. *Biochim. Biophys. Acta* **679**:116–124.
  68. **Krinner, M., G. Hauska, E. Hurt, and W. Lockau.** 1982. A cytochrome *f-b<sub>6</sub>* complex with plastoquinol-cytochrome c oxidoreductase activity from *Anabaena variabilis*. *Biochim. Biophys. Acta* **681**:110–117.
  69. **Lange, O. L., W. Bilger, S. Rimke, and U. Schreiber.** 1989. Chlorophyll fluorescence of lichens containing green and blue-green algae during hydration by water vapor uptake and by addition of liquid water. *Bot. Acta* **102**:306–313.
  70. **Laudenbach, D. E., and N. A. Straus.** 1988. Characterization of a cyanobacterial iron stress-induced gene similar to *psbC*. *J. Bacteriol.* **170**:5018–5026.
  71. **Laudenbach, D. E., S. K. Herbert, C. McDowell, D. C. Fork, A. Grossman, and N. A. Straus.** 1990. Cytochrome c-553 is not required for photosyn-

- thetic activity in the cyanobacterium *Synechococcus*. *Plant Cell* **2**:913–924.
72. Lee, T., M. Tsuzuki, K. Yokoyama, and I. Karube. 1994. *In-Vivo* fluorometric method for early detection of cyanobacterial waterblooms. *J. App. Phycol.* **6**:489–495.
  73. Leisner, J. M. R., W. Bilger, and O. L. Lange. 1996. Chlorophyll fluorescence characteristics of the cyanobacterial lichen *Peltigera rufescens* under field conditions. I. Seasonal patterns of photochemical activity and the occurrence of Photosystem II inhibition. *Flora* **191**:261–273.
  74. Leisner, J. M. R., T. G. A. Green, and O. L. Lange. 1997. Photobiont activity of a temperate crustose lichen—long-term chlorophyll fluorescence and CO<sub>2</sub> exchange measurements in the field. *Symbiosis* **23**:165–182.
  75. Leonhardt, K. G., and N. A. Straus. 1992. An iron stress operon involved in photosynthetic electron transport in the marine cyanobacterium *Synechococcus* sp. PCC 7002. *J. Gen. Microbiol.* **138**:1613–1621.
  76. Leonhardt, K. G., and N. A. Straus. 1994. Photosystem II genes *isiA*, *psbD1* and *psbC* in *Anabaena* sp. PCC 7120: cloning, sequencing and the transcriptional regulation in iron-stressed and iron-repleted cells. *Plant Mol. Biol.* **24**:63–73.
  77. Li, Q. L., and D. T. Canvin. 1997. Inorganic carbon accumulation stimulates linear electron flow to artificial electron-acceptors of Photosystem-I in air-grown cells of the cyanobacterium *Synechococcus* UTEX-625. *Plant Physiol.* **114**:1273–1281.
  78. Li, Q. L., and D. T. Canvin. 1997. Effect of the intracellular inorganic carbon pool on chlorophyll-a fluorescence quenching and O<sub>2</sub> photoreduction in air-grown cells of the cyanobacterium *Synechococcus* UTEX-625. *Can. J. Bot.* **75**:946–954.
  79. Lien, S., and A. San Pietro. 1979. On the reactivity of oxygen with photosystem I electron acceptors. *FEBS Lett.* **99**:189–193.
  80. Lüttge, U., B. Büdel, E. Ball, F. Strube, and P. Weber. 1995. Photosynthesis of terrestrial cyanobacteria under light and desiccation stress as expressed by chlorophyll fluorescence and gas exchange. *J. Exp. Bot.* **46**:309–319.
  81. Manna, P., and W. Vermass. 1997. Lumenal proteins involved in respiratory electron-transport in the cyanobacterium *Synechocystis* sp. PCC 6803. *Plant Mol. Biol.* **35**:407–416.
  82. Maxwell, D. P., S. Falk, and N. P. A. Huner. 1995. Photosystem II excitation and development of resistance to photoinhibition. *Plant Physiol.* **107**:687–694.
  83. Meunier, P. C., J. W. Watters, M. S. Colón-López, and L. A. Sherman. 1995. Regulation of the O<sub>2</sub>-evolving mechanism during N<sub>2</sub> fixation in the diazotrophic cyanobacterium *Cyanothece* sp. ATCC 51142, p. 389–392. *In P. Mathis* (ed.), *Research in photosynthesis*, vol. II, Kluwer Academic Publishers, Dordrecht, The Netherlands.
  84. Meunier, P. C., R. L. Burnap, and L. A. Sherman. 1995. Interaction of the photosynthetic and respiratory electron transport chains producing slow O<sub>2</sub> signals under flashing light in *Synechocystis* sp. PCC 6803. *Photosynth. Res.* **45**:31–40.
  85. Mi, H. L., T. Endo, U. Schreiber, T. Ogawa, and K. Asada. 1992. Electron donation from cyclic and respiratory flows to the photosynthetic intersystem chain is mediated by pyridine nucleotide dehydrogenase in the cyanobacterium *Synechocystis* sp. PCC 6803. *Plant Cell Physiol.* **33**:1233–1237.
  86. Mi, H. L., T. Endo, T. Ogawa, and K. Asada. 1995. Thylakoid membrane-bound, NADPH-specific pyridine nucleotide dehydrogenase complex mediates cyclic electron transport in the cyanobacterium *Synechocystis* sp. PCC 6803. *Plant Cell Physiol.* **36**:661–668.
  - 86a. Miller, A. Personal communication.
  87. Miller, A. G., and D. T. Canvin. 1987. The quenching of chlorophyll a fluorescence as a consequence of the transport of inorganic carbon by the cyanobacterium *Synechococcus* UTEX 625. *Biochim. Biophys. Acta* **894**:407–413.
  88. Miller, A. G., G. S. Espie, and D. T. Canvin. 1990. Active transport of inorganic carbon increases the rate of O<sub>2</sub> photo-reduction by the cyanobacterium *Synechococcus* UTEX 625. *Plant Physiol.* **88**:6–9.
  89. Miller, A. G., G. S. Espie, and D. Bruce. 1996. Characterization of the non-photochemical quenching of chlorophyll fluorescence that occurs during the active accumulation of inorganic carbon in the cyanobacterium *Synechococcus* PCC 7942. *Photosynth. Res.* **49**:251–262.
  90. Miller, A. G., G. S. Espie, and D. T. Canvin. 1988. Chlorophyll a fluorescence yield as a monitor of both active CO<sub>2</sub> and HCO<sub>3</sub><sup>-</sup> transport by the cyanobacterium *Synechococcus* UTEX 6251. *Plant Physiol.* **86**:655–658.
  91. Miller, A. G., G. S. Espie, and D. T. Canvin. 1991. The effects of inorganic carbon and oxygen on fluorescence in the cyanobacterium *Synechococcus* UTEX 625. *Can. J. Bot.* **69**:1151–1160.
  92. Mir, N. A., C. Salom, and D. T. Canvin. 1995. Photosynthetic nitrite reduction as influenced by the internal inorganic carbon pool in air-grown cells of *Synechococcus* UTEX-625. *Plant Physiol.* **108**:313–318.
  93. Miyachi, S., K. Kotzabasis, J. Thielmann, H. Senger, and J. Burger. 1996. Photosynthetic characteristics of 3 strains of cyanobacteria grown under low-CO<sub>2</sub> or high-CO<sub>2</sub> conditions. *Z. Naturforsch. Sect. C* **51**:40–46.
  94. Mullineaux, C. W., and J. F. Allen. 1986. The state 2 transition in the cyanobacterium *Synechococcus* 6301 can be driven by respiratory electron flow into the plastoquinone pool. *FEBS Lett.* **205**:155–160.
  95. Mullineaux, C. W., and J. F. Allen. 1990. State 1-state 2 transitions in the cyanobacterium *Synechococcus* 6301 are controlled by the redox state of electron carriers between Photosystems I and II. *Photosynth. Res.* **23**:297–311.
  96. Mullineaux, C. W., M. J. Tobin, and G. R. Jones. 1997. Mobility of photosynthetic complexes in thylakoid membranes. *Nature* **390**:421–424.
  97. Murata, N. 1969. Control of excitation transfer in photosynthesis. I. Light-induced change of chlorophyll a fluorescence in *Porphyridium cruentum*. *Biochim. Biophys. Acta* **172**:242–251.
  98. Myers, J., J. R. Graham, and R. T. Wang. 1980. Light harvesting in *Anacystis nidulans* studied in pigment mutants. *Plant Physiol.* **66**:1144–1149.
  99. Ögren, E. 1994. The significance of photoinhibition for photosynthetic productivity, p. 433–447. *In N. R. Baker and J. R. Bowyer* (ed.), *Photoinhibition of photosynthesis: from molecular mechanisms to the field*. Bios Scientific Publishers, Oxford, United Kingdom.
  100. Ögren, E., and G. Öquist. 1984. Photoinhibition of photosynthesis in *Lemna gibba* as induced by the interaction between light and temperature. II. Photosynthetic electron transport. *Physiol. Plant.* **62**:187–192.
  101. Ögren, E., and E. Rosenqvist. 1992. On the significance of photoinhibition of photosynthesis in the field and its generality among species. *Photosynth. Res.* **33**:63–71.
  102. Öquist, G., J. M. Anderson, S. McCaffery, and W. S. Chow. 1992. Mechanistic differences in photoinhibition of sun and shade plants. *Planta* **188**:422–431.
  103. Öquist, G., and N. P. A. Huner. 1993. Cold-hardening-induced resistance to photoinhibition of photosynthesis in winter rye is dependent upon an increased capacity for photosynthesis. *Planta* **189**:150–156.
  104. Öquist, G., and W. S. Chow. 1992. On the relationship between the quantum yield of Photosystem II electron transport, as determined by chlorophyll fluorescence and the quantum yield of CO<sub>2</sub>-dependent O<sub>2</sub> evolution. *Photosynth. Res.* **33**:51–62.
  105. Öquist, G., D. Campbell, A. K. Clarke, and P. Gustafsson. 1995. The cyanobacterium *Synechococcus* modulates PS II function in response to excitation stress through D1 exchange. *Photosynth. Res.* **46**:151–158.
  106. Öquist, G., V. M. Hurry, and N. P. A. Huner. 1993. Low-temperature effects on photosynthesis and correlation with freezing tolerance in spring and winter cultivars of wheat and rye. *Plant Physiol.* **101**:245–250.
  107. Öquist, G., V. M. Hurry, and N. P. A. Huner. 1993. The temperature dependence of the redox state of QA and susceptibility of photosynthesis to photoinhibition. *Plant Physiol. Biochem.* **31**:683–691.
  108. Palmqvist, K. 1995. Uptake and fixation of CO<sub>2</sub> in lichen photobionts. *Symbiosis* **18**:95–109.
  109. Palmqvist, K. 1993. Photosynthetic CO<sub>2</sub>-use efficiency in lichens and their isolated photobionts: the possible role of a CO<sub>2</sub>-concentrating mechanism. *Planta* **191**:48–56.
  110. Palmqvist, K., G. Samuelsson, and M. R. Badger. 1994. Photobiont-related differences in carbon acquisition among green-algal lichens. *Planta* **195**:70–79.
  111. Papageorgiou, G., and Govindjee. 1968. Light-induced changes in the fluorescence yield of chlorophyll a in vivo. 1. *Anacystis nidulans*. *Biophys. J.* **8**:1299–1315.
  112. Papageorgiou, G. C. 1996. The photosynthesis of cyanobacteria (blue bacteria) from the perspective of signal analysis of chlorophyll a fluorescence. *J. Sci. Ind. Res.* **55**:596–617.
  - 112a. Park, Y.-I., G. Samuelsson, and G. Öquist. Unpublished data.
  113. Peschek, G. A., and G. Schmetterer. 1982. Evidence for plastoquinol-cytochrome *fb*-563 oxidoreductase as a common electron donor to P-700 and cytochrome oxidase in cyanobacteria. *Biochem. Biophys. Res. Commun.* **108**:1188–1195.
  114. Rippka, R., J. Deruelles, J. B. Waterbury, M. Herdman, and R. Y. Stanier. 1979. Generic assignments, strain histories and properties of pure cultures of cyanobacteria. *J. Gen. Microbiol.* **111**:1–61.
  115. Rippka, R., J. Waterbury, and G. Cohen-Bazire. 1974. A cyanobacterium which lacks thylakoids. *Arch. Microbiol.* **100**:419–436.
  116. Romero, J. M., C. Lara, and M. N. Sivak. 1992. Effect of carbon and nitrogen assimilation on chlorophyll fluorescence emission by the cyanobacterium *Anacystis nidulans*. *Physiol. Plant.* **85**:433–438.
  117. Sabbert, D., S. Engelbrecht, and W. Junge. 1996. Intersubunit rotation in active F-ATPase. *Nature* **381**:623–625.
  118. Salehian, O., and D. Bruce. 1992. Distribution of excitation energy in photosynthesis, quantification of fluorescence yields from intact cyanobacteria. *J. Lumin.* **51**:91–98.
  119. Sandmann, G. 1986. Formation of plastocyanin and cytochrome c-553 in different species of blue-green algae. *Arch. Microbiol.* **145**:76–79.
  120. Sandmann, G., and P. Böger. 1980. Copper induced exchange of plastocyanin and cytochrome c-553 in cultures of *Anabaena variabilis* and *Plectonema borianum*. *Plant Sci.* **17**:417–424.
  121. Scherer, S., E. Stürzl, and P. Böger. 1982. Interaction of respiratory and photosynthetic electron transport in *Anabaena variabilis* Kütz. *Arch. Microbiol.* **132**:333–337.
  122. Scherer, S., H. Almon, and P. Böger. 1988. Interaction of photosynthesis, respiration and nitrogen fixation in cyanobacteria. *Photosynth. Res.* **15**:95–114.

123. Schneegurt, M., D. M. Sherman, S. Nayar, and L. M. Sherman. 1994. Oscillating behaviour of carbohydrate granule formation and dinitrogen fixation in the cyanobacterium *Cyanothece* sp. strain ATCC 51142. *J. Bacteriol.* **176**:1586–1597.
124. Schneegurt, M. A., D. M. Sherman, and L. M. Sherman. 1997. Composition of the carbohydrate granules of the cyanobacterium, *Cyanothece* sp. strain ATCC 51142. *Arch. Microbiol.* **167**:89–98.
125. Schreiber, U. 1994. New emitter-detector-cuvette assembly for measuring modulated chlorophyll fluorescence of highly diluted suspensions in conjunction with the standard PAM fluorometer. *Z. Naturforsch. C* **49**:646–656.
126. Schreiber, U., R. Gademann, P. J. Ralph, and A. W. D. Larkum. 1997. Assessment of photosynthetic performance of *Prochloron* in *Lissoclinum patella* in hospite by chlorophyll fluorescence measurements. *Plant Cell Physiol.* **38**:945–951.
127. Schreiber, U., H. Hormann, C. Neubauer, and C. Klughammer. 1995. Assessment of photosystem II photochemical quantum yield by chlorophyll fluorescence quenching analysis. *Aust. J. Plant Physiol.* **22**:209–220.
128. Schreiber, U., T. Endo, H. Mi, and K. Asada. 1995. Quenching analysis of chlorophyll fluorescence by the saturation pulse method: particular aspects relating to the study of eukaryotic algae and cyanobacteria. *Plant Cell Physiol.* **36**:873–882.
129. Schreiber, U., U. Schliwa, and W. Bilger. 1986. Continuous recording of photochemical and non-photochemical chlorophyll fluorescence quenching with a new type of modulation fluorometer. *Photosynth. Res.* **10**:51–62.
130. Schroeter, B. 1994. *In-situ* photosynthetic differentiation of the green algal and the cyanobacterial photobiont in the crustose lichen *Placopsis contortuplicata*. *Oecologia* **98**:212–220.
131. Schubert, H., H. Matthijs, and L. Mur. 1995. In vivo assay of P700 redox changes in the cyanobacterium *Fremyella diplosiphon* and the role of cytochrome-c-oxidase in regulation of photosynthetic electron transfer. *Photosynthetica* **31**:517–527.
132. Schubert, H., H. C. P. Matthijs, L. R. Mur, and U. Schiewer. 1995. Blooming of cyanobacteria in turbulent water with steep light gradients: the effect of intermittent light and dark periods on the oxygen evolution capacity of *Synechocystis* sp. PCC 6803. *FEMS Microbiol. Ecol.* **18**:237–245.
133. Schubert, H., R. Forster, and S. Sager. 1995. In situ measurement of state transition in cyanobacterial blooms: kinetics and extent of the state change in relation to underwater light and vertical mixing. *Mar. Ecol. Prog. Ser.* **128**:99–108.
134. Seaton, G. R., and D. A. Walker. 1990. Chlorophyll fluorescence as a measure of photosynthetic carbon metabolism. *Proc. R. Soc. London Ser. B* **242**:29–35.
135. Shyam, R., A. S. Raghavendra, and P. V. Sane. 1993. Role of dark respiration in photoinhibition of photosynthesis and its reactivation in the cyanobacterium *Anacystis nidulans*. *Physiol. Plant.* **88**:446–452.
136. Sidler, W. 1994. Phycobilisome and phycobiliprotein structures, p. 139–216. In D. A. Bryant (ed.), *The molecular biology of cyanobacteria*. Kluwer Academic Publishers, Dordrecht, The Netherlands.
137. Strasser, R. J., A. Srivastava, and Govindjee. 1995. Polyphasic chlorophyll-alpha fluorescence transient in plants and cyanobacteria. *Photochem. Photobiol.* **61**:32–42.
138. Sundberg, B., D. Campbell, and K. Palmqvist. 1997. Predicting CO<sub>2</sub> gain and photosynthetic light acclimation from fluorescence yield and quenching in cyano-lichens. *Planta* **201**:138–145.
139. Tanaka, Y., S. Katada, H. Ishikawa, T. Ogawa, and T. Takabe. 1997. Electron flow from NAD(P)H dehydrogenase to Photosystem-I is required for adaptation to salt shock in the cyanobacterium-*Synechocystis* sp. PCC 6803. *Plant Cell Physiol.* **38**:1311–1318.
140. Tandeau de Marsac, N., and J. Houmard. 1988. Complementary chromatic adaptation: physiological conditions and action spectra. *Methods Enzymol.* **167**:318–328.
141. Ting, C. S., and T. G. Owens. 1992. Limitations of the pulse-modulated technique for measuring the fluorescence characteristics of algae. *Plant Physiol.* **100**:367–373.
142. Torzillo, G., P. Accolla, E. Pinzani, and J. Masojidek. 1996. *In-situ* monitoring of chlorophyll fluorescence to assess the synergistic effect of low-temperature and high irradiance stresses in *Spirulina* cultures grown outdoors in photobioreactors. *J. Appl. Phycol.* **8**:283–291.
143. Trebst, A. 1980. Inhibitors in electron flow. *Methods Enzymol.* **69**:675–715.
144. Trissl, H.-W., and J. Lavergne. 1995. Fluorescence induction from photosystem II: analytical equations for the yields of photochemistry and fluorescence derived from analysis of a model including exciton-radical pair equilibrium and restricted energy transfer between photosynthetic units. *Aust. J. Plant Physiol.* **22**:183–193.
145. van Kooten, O., and J. F. H. Snel. 1990. The use of chlorophyll fluorescence nomenclature in plant stress physiology. *Photosynth. Res.* **25**:147–150.
146. van Walraven, H. S., H. Strotman, O. Schwarz, and B. Rumberg. 1996. The H<sup>+</sup>/ATP coupling ratio of the ATP synthase from thiol-modulated chloroplasts and two cyanobacterial strains is four. *FEBS Lett.* **379**:309–313.
147. Vermaas, W. F. J., G. Shen, and S. Styring. 1994. Electrons generated by photosystem II are utilized by an oxidase in the absence of photosystem I in the cyanobacterium *Synechocystis* sp. PCC 6803. *FEBS Lett.* **337**:103–108.
148. Vernotte, C., A. L. Etienne, and J. M. Briantais. 1979. Quenching of the system II chlorophyll fluorescence by the plastoquinone pool. *Biochim. Biophys. Acta* **545**:519–527.
149. Vernotte, C., C. Astier, and J. Olive. 1990. State 1-state 2 adaptation in the cyanobacteria *Synechocystis* PCC 6714 wild type and *Synechocystis* PCC 6803 wild type and phycocyanin-less mutant. *Photosynth. Res.* **26**:203–212.
150. Vonshak, A., G. Torzillo, P. Accolla, and L. Tomaselli. 1996. Light and oxygen stress in *Spirulina platensis* (cyanobacteria) grown outdoors in tubular reactors. *Physiol. Plant.* **97**:175–179.
151. Walker, D. 1987. The use of the oxygen electrode and fluorescence probes in simple measurements of photosynthesis. Hansatech Instruments Ltd., Kings Lynn, United Kingdom.
152. Walters, R. G., and P. Horton. 1993. Theoretical assessment of alternative mechanisms for non-photochemical quenching of PS II fluorescence in barley leaves. *Photosynth. Res.* **36**:119–139.
153. Weis, E., and J. A. Berry. 1987. Quantum efficiency of photosystem II in relation to energy dependent quenching of chlorophyll fluorescence. *Biochim. Biophys. Acta* **894**:198–208.
154. Zhang, L., B. McSpadden, H. B. Pakarasi, and J. Whitmarsh. 1992. Copper mediated regulation of cytochrome c553 and plastocyanin in the cyanobacterium *Synechocystis* 6803. *J. Biol. Chem.* **267**:19054–19059.
155. Zhang, L., H. B. Pakarasi, and J. Whitmarsh. 1994. Photoautotrophic growth of the cyanobacterium *Synechocystis* sp. PCC 6803 in the absence of cytochrome c<sub>553</sub> and plastocyanin. *J. Biol. Chem.* **269**:5036–5042.
156. Zhao, J., J. Zhou, and D. A. Bryant. 1992. Energy transfer processes in phycobilisomes as deduced from analyses of mutants of *Synechococcus* sp. PCC 7002, p. 25–32. In N. Murata (ed.) *Research in photosynthesis*, vol 1. Kluwer Academic Publishers, Dordrecht, The Netherlands.
157. Zhou, G., and P. Gustafsson. Unpublished data.
158. Zhou, G. Unpublished data.