

## *Helicobacter pylori* Physiology Predicted from Genomic Comparison of Two Strains

PETER DOIG,<sup>1\*</sup> BOUDEWIJN L. DE JONGE,<sup>1</sup> RICHARD A. ALM,<sup>1</sup> ERIC D. BROWN,<sup>1†</sup> MARIA URIA-NICKELSEN,<sup>1</sup>  
BRIAN NOONAN,<sup>1</sup> SCOTT D. MILLS,<sup>1</sup> PETER TUMMINO,<sup>1</sup> GILLES CARMEL,<sup>2</sup> BRAYDON C. GUILD,<sup>2</sup>  
DONALD T. MOIR,<sup>2‡</sup> GERALD F. VOVIS,<sup>2‡</sup> AND TREVOR J. TRUST<sup>1</sup>

*AstraZeneca R&D Boston, Cambridge,<sup>1</sup> and Genome Therapeutics Corp., Waltham,<sup>2</sup> Massachusetts*

INTRODUCTION .....	675
ANALYSES OF GENETIC AND FUNCTIONAL CONSERVATION .....	676
NUTRITIONAL REQUIREMENTS .....	676
Amino Acids and Polyamines .....	676
Cofactors and Vitamins .....	692
Purine and Pyrimidine Biosynthesis, Salvage, and Interconversion .....	692
Inorganic Elements and Heavy Metals .....	695
Carbohydrates .....	697
CENTRAL INTERMEDIARY AND ENERGY METABOLISM .....	697
Central Intermediary Metabolism .....	697
Glycolysis and gluconeogenesis .....	697
Entner-Doudoroff and phosphopentose pathways .....	697
Pyruvate metabolism .....	697
Fermentation .....	698
Tricarboxylic acid cycle .....	698
Fatty acid degradation .....	698
Electron Transport Chain .....	698
Electron donors .....	698
Quinones and cytochromes .....	698
Terminal electron acceptors .....	698
ATP-Proton Motive Force Conversion .....	698
Detoxification .....	699
MACROMOLECULE BIOSYNTHESIS AND MODIFICATION .....	699
DNA Replication, Recombination, and Restriction-Modification .....	699
Transcription and Translation .....	699
Fatty Acid and Phospholipid .....	700
Peptidoglycan .....	700
Outer Membrane .....	701
Lipopolysaccharide .....	701
Flagella .....	701
CELLULAR PROCESSES .....	701
Protein Secretion .....	701
Cag Pathogenicity Island .....	702
Insertion Elements .....	702
Transformation .....	702
Chemotaxis .....	702
Cell Division and Morphology .....	703
Virulence Factors .....	703
CONCLUSION .....	703
REFERENCES .....	704

### INTRODUCTION

*Helicobacter pylori* is a gram-negative bacterium which colonizes the gastric mucosa of humans, causes gastritis and pep-

tic ulcer disease, and is associated with certain types of gastric cancer (27, 65, 87). Once colonized, the host can be chronically infected for life unless antimicrobial therapy is administered. The ability to colonize and persist in the human stomach for many years indicates that *H. pylori* is specifically adapted to occupy only this niche, and such adaptation should be reflected in a unique complement of physiological capabilities. Furthermore, physiological differences resulting from the apparent genomic variation among strains have been suggested to be responsible for the diversity of diseases associated with *H. pylori* infection (16, 95).

Bacterial genomics, the identification and annotation of the

\* Corresponding author. Mailing address: AstraZeneca R&D Boston, 128 Sidney St., Cambridge, MA 02139. Phone: (617) 234-2534. Fax: (617) 576-3030. E-mail: Peter.Doig@arcb.us.astra.com.

† Present address: Department of Biochemistry, Health Sciences, McMaster University, Hamilton, Ontario, Canada, L8N 3Z5.

‡ Present address: Genaissance Pharmaceuticals, New Haven, CT 06511.

entire coding potential of a bacterium, allows a more complete understanding of bacterial physiology and pathogenesis. The recent analysis of the complete genomic sequence of two unrelated, pathogenic *H. pylori* strains (J99 and 26695) demonstrated that even though the chromosomes are organized differently in a limited number of discrete regions, the genome size, genetic content, and gene order of these two strains are remarkably similar (6). We have used the data resulting from this comparative sequence analysis as the starting point to examine, from a functional perspective, the genes that are common and unique to the two strains. The presence or absence of orthologous genes or metabolic pathways in both unrelated *H. pylori* strains implies that these genes or pathways are present or absent, respectively, in this species. Our comparison has defined the set of common *H. pylori* metabolic capabilities as well as a small number that are strain specific.

### ANALYSES OF GENETIC AND FUNCTIONAL CONSERVATION

The predicted genes from both *H. pylori* genomes (6) were assigned a likely function, if their predicted amino acid sequence exhibited similarity to a protein of known function, and categorized as shown in Table 1. Function was annotated conservatively; e.g., proteins which showed sequence similarity to transporters, the majority of which did not have the same substrate specificity, were assigned to the general category of transporters pending experimental evidence for their specificity. The two *H. pylori* genomes are highly conserved with respect to gene content (1,495 and 1,552 open reading frames [ORFs] in J99 and 26695, respectively [6]), functional categorization (Table 2), and gene order (Fig. 1). In both strains, approximately 58% of the gene products were assigned a putative function based upon their having significant sequence similarity to a protein of known function; nearly 18% were conserved in other species but had no known function; and about 23% were specific to *H. pylori* (Table 2). Eighty-nine genes were specific to strain J99, and 117 were specific to strain 26695; 26 of these genes in each of the strains J99 and 26695 had an assigned function. Preliminary analysis of the *Campylobacter jejuni* genome (based on analysis of the recently completed genome by the Sanger Centre) indicated that approximately 90 of the *H. pylori* specific genes will have an orthologue in this closely related species. This will reduce the proportion of *H. pylori*-specific genes to approximately 17%.

Comparison of orthologous genes and their encoded products showed a high degree of conservation. Sequence variation between the two strains was significantly greater at the nucleotide level than at the amino acid level. Because the nucleotide variation occurred most commonly in the third position of a coding triplet, the primary sequence of the encoded protein was highly conserved (Table 3). The fact that many of the nucleotide differences are silent with respect to the protein sequence suggests that there is a strong selective pressure for functional conservation at the protein level.

The nucleotide drift in the third position of a coding triplet is probably responsible for the majority of the DNA-based "diversity" reported for *H. pylori* (3, 4, 12, 54, 75, 77, 148). For example, pulsed-field gel electrophoresis mapping data have been interpreted to mean that the gene order and physical arrangement of the chromosome are highly variable from strain to strain (75, 148). By using this technique, strain J99 and strain 26695 would appear to be highly divergent in both the number of *NotI* fragments and gene location (6). This apparent genetic diversity is easily explained by two inversions in combination with the silent nucleotide drift, which is responsible

for six of the seven additional *NotI* sites found in strain J99 compared to strain 26695 (6). Although the genomic content and the resulting physiological capabilities of the two strains are almost identical, these few differences in gene arrangement would have classified these strains as diverse. This example reveals the limitations of DNA-based methods when used to examine strain diversity.

## NUTRITIONAL REQUIREMENTS

### Amino Acids and Polyamines

Both sequenced strains of *H. pylori* have homologues to all the genes that would be needed to synthesize eight amino acids from central intermediary metabolites (Table 4). Studies of the growth requirements for several strains of *H. pylori* have shown an absolute need for arginine, histidine, leucine, isoleucine, valine, methionine, and phenylalanine (118, 132), a finding consistent with the genomic sequence information.

Although no homologues to the genes involved in the amidation of aspartate were detected in *H. pylori*, several strains have been reported to grow in the absence of asparagine (132). It is possible that asparagine is synthesized by an aspartyl-tRNA-asparagine amidotransferase, similar to what has been observed with glutamyl-tRNA biosynthesis in *Bacillus subtilis* (see "Transcription and translation" below) (29).

Both the serine and tyrosine biosynthetic pathways were complete except for a homologue to their respective specific transaminase. However, each of these reactions may be catalyzed by one of the several identified transaminases with undetermined substrate specificity (JHP206/HP220, JHP568/HP624, JHP673/HP736, and JHP976/HP405). Such an enzymatic activity would allow the de novo synthesis of these amino acids, as observed in some strains of *H. pylori* (132). Regardless of whether *H. pylori* can synthesize serine, it possesses a specific transporter which allows the acquisition of this amino acid from the environment. *sdaC*, which encodes the serine transporter, is contiguous with *sdaB*, whose protein product in *Escherichia coli* converts L-serine to pyruvate. Similarly, *putP*, which encodes a proline transporter, is adjacent to *putA*, which encodes a bifunctional enzyme that oxidizes proline to L-glutamate in *E. coli*. The alanine transporter gene (JHP877/HP0942) is clustered with two other genes (*alr* and *dadA*) which are involved in alanine metabolism (Fig. 2). A positive regulator of the *dad* operon is thought to be upstream of this gene cluster in *E. coli*. In *H. pylori*, a putative regulatory gene (JHP879/HP0944) has also been identified upstream of the *dad* gene cluster. This putative regulatory gene does not have homology to the putative *E. coli* regulator, a finding which may indicate that the regulation of alanine catabolism is different in these two species. The *H. pylori* gene clusters described above would allow for the uptake and utilization of serine, proline, and alanine as carbon and nitrogen sources. In addition, *H. pylori* has a transporter for the uptake of glutamate (JHP1399/HP1506), an amino acid abundant in gastric juice (82).

*H. pylori* also encodes homologues for four other amino acid uptake systems with unknown specificity. One of these systems consists of a gene cluster encoding a multisubunit periplasmic permease (JHP1096–1099/HP1169–1172). The putative operon encodes two permeases, an ATP-binding protein and a periplasmic binding protein. Based on sequence similarity, the ligand for this high-affinity transporter may be either glutamine, histidine, or arginine. *H. pylori* is unable to synthesize the last two amino acids and therefore requires transport systems for them. Furthermore, the apparent inability of *H. pylori* to synthesize

TABLE 1. List of *H. pylori* J99 genes and corresponding 26695 orthologs with putative functional assignments<sup>a</sup>

Gene no. in:		Gene name	Function
J99	26695		
<b>Amino acid biosynthesis</b>			
General			
206	0220		Aminotransferase
568	0624		Aminotransferase
632	0696		Hydantoin utilization
633	0695		Hydantoin utilization
673	0736		Aminotransferase
976	0405		Aminotransferase
Aromatic amino acid family			
122	0134	<i>aroF</i>	Phospho-2-dehydro-3-deoxyheptonate aldolase
145	0157	<i>aroK</i>	Shikamate kinase I
268	0283	<i>aroB</i>	3-Dehydroquinate synthase
386	1038	<i>aroD</i>	3-Dehydroquinate dehydratase
608	0663	<i>aroC</i>	Chorismate synthase
980	0401	<i>aroA</i>	3-Phosphoshikimate 1-carboxyvinyl transferase
1170	1249	<i>aroE</i>	Shikimate 5-dehydrogenase
1198	1277	<i>trpA</i>	Tryptophan synthase alpha chain
1199	1278	<i>trpB</i>	Tryptophan synthase beta chain
1200	1279	<i>trpC</i>	Indole-3-glycerol phosphate synthase
1201	1280	<i>trpD</i>	Anthranilate phosphoribosyltransferase
1202	1281	<i>trpG</i>	Anthranilate synthase component II
1203	1282	<i>trpE</i>	Anthranilate synthase component I
1294	1380	<i>tyrA</i>	Prephenate dehydrogenase
Aspartate family			
90	0098	<i>thrC</i>	Threonine synthase
98	0106	<i>metB</i>	Cystathionine gamma-synthase
198	0212	<i>dapE</i>	Succinyl-diaminopimelate desuccinylase
275	0290	<i>lysA</i>	Diaminopimelate decarboxylase
375	1050	<i>thrB</i>	Homoserine kinase
410	1013	<i>dapA</i>	Dihydrodipicolinate synthase
460	0510	<i>dapB</i>	Dihydrodipicolinate reductase
513	0566	<i>dapF</i>	Diaminopimelate epimerase
570	0626	<i>dapD</i>	2,3,4,5-Tetrahydropyridine-2-carboxylate- <i>N</i> -succinyltransferase
594	0649	<i>aspA</i>	Aspartate ammonia-lyase
615	0672	<i>aspB</i>	Aspartate aminotransferase
761	0822	<i>hom</i>	Homoserine dehydrogenase
1114	1189	<i>asd</i>	Aspartate-semialdehyde dehydrogenase
1150	1229	<i>lysC</i>	Aspartokinase 2 alpha and beta subunits
Glutamate family			
461	0512	<i>glnA</i>	Glutamine synthetase
1001	0380	<i>gdhA</i>	Glutamate dehydrogenase
1085	1158	<i>proC</i>	Pyrroline-5-carboxylate reductase
Pyruvate family			
313	0330	<i>ilvC</i>	Ketol-acid reductoisomerase
1361	1468	<i>ilvE</i>	Branched-chain amino acid aminotransferase
Serine family			
99	0107	<i>cysK</i>	Cysteine synthase
171	0183	<i>glyA</i>	Serine hydroxymethyltransferase
597	0652	<i>serB</i>	Phosphoserine phosphatase
984	0397	<i>serA</i>	Phosphoglycerate dehydrogenase
1133	1210	<i>cysE</i>	Serine acetyltransferase
<b>Biosynthesis of cofactors, prosthetic groups, and carriers</b>			
Biotin			
25	0029	<i>bioD</i>	Dethiobiotin synthetase
545	0598	<i>bioF</i>	8-Amino-7-oxononanoate synthase
910	0976	<i>bioA</i>	Adenosylmethionine-8-amino-7-oxononanoate aminotransferase
1068	1140		Biotin activation protein
1298	1406	<i>bioB</i>	Biotin synthetase
Folic acid			
278	0293	<i>pabB</i>	<i>p</i> -Aminobenzoate synthetase
388	1036	<i>folK</i>	7,8-Dihydro-6-hydroxymethylpterin-pyrophosphokinase
524	0577	<i>folD</i>	Methylenetetrahydrofolate dehydrogenase/methenyltetrahydrofolate cyclohydrolase
862	0928	<i>folE</i>	GTP cyclohydrolase I

Continued on following page

TABLE 1—Continued

Gene no. in:		Gene name	Function
J99	26695		
863	0928	<i>folE</i>	GTP cyclohydrolase I
1153	1232	<i>folP</i>	Dihydropteroate synthase
1403	1510	<i>folB</i>	Dihydroneopterin aldolase
1454	1545	<i>folC</i>	Folylpolylglutamate synthase
Heme and porphyrin			
150	0163	<i>hemB</i>	δ-Aminolevulinic acid dehydratase
222	0237	<i>hemC</i>	Porphobilinogen deaminase
224	0239	<i>hemA</i>	Glutamyl-tRNA reductase
291	0306	<i>hemL</i>	Glutamate-1-semialdehyde-2,1-aminomutase
551	0604	<i>hemE</i>	Uroporphyrinogen decarboxylase
610	0665	<i>hemN</i>	Oxygen-independent coproporphyrinogen III oxidase
1000	0381	<i>hemG</i>	Protoporphyrinogen oxidase
1005	0376	<i>hemH</i>	Ferrochelataase
1145	1224	<i>hemD</i>	Uroporphyrinogen III synthase
1147	1226	<i>hemN</i>	Oxygen-independent coproporphyrinogen III oxidase
Menaquinone and ubiquinone			
225	0240	<i>ispB</i>	Octaprenyl-diphosphate synthase
864	0929	<i>ispA</i>	Geranyltransferase
1278	1360	<i>ubiA</i>	4-Hydroxybenzoate octaprenyltransferase
1369	1476	<i>ubiD</i>	Octaprenyl-4-hydroxybenzoate carboxy-lyase
1376	1483	<i>ubiE</i>	Ubiquinone/menaquinone biosynthesis methyltransferase
Molybdopterin			
158	0172	<i>moeA</i>	Molybdopterin biosynthesis protein
705	0768	<i>moaA</i>	Molybdopterin cofactor biosynthetic protein
706	0769	<i>mobA</i>	Molybdopterin-guanine dinucleotide biosynthesis protein A
734	0798	<i>moaC</i>	Molybdenum cofactor biosynthesis protein C
735	0799	<i>mog</i>	Molybdopterin biosynthesis protein
736	0800	<i>moaE</i>	Molybdopterin-converting factor, subunit 2
737	0801	<i>moaD</i>	Molybdopterin-converting factor, subunit 1
750	0814	<i>moeB</i>	Molybdopterin-synthase sulfurlyase
Pantothenate			
6	0006	<i>panC</i>	Pantoate-β-alanine ligase
30	0034	<i>panD</i>	Aspartate-1-decarboxylase
367	1058	<i>panB</i>	3-Methyl-2-oxobutanoate hydroxymethyltransferase
779	0841	<i>dfp</i>	Pantothenate metabolism flavoprotein
Pyridoxine			
328	0354	<i>dxs</i>	1-Deoxyxylulose-5-phosphate synthase
1489	1582	<i>pdxJ</i>	Pyridoxal phosphate synthetase
1490	1583	<i>pdxA</i>	Pyridoxal phosphate biosynthetic protein A
Riboflavin			
2	0002	<i>ribE</i>	Riboflavin synthase beta chain
338	1087	<i>ribF</i>	Riboflavin kinase
738	0802	<i>ribA</i>	GTP cyclohydrolase II
740	0804	<i>ribBA</i>	GTP cyclohydrolase II/3,4-dihydroxy-2-butanone 4-phosphate synthase
1398	1505	<i>ribD</i>	Riboflavin-specific deaminase
1482	1574	<i>ribC</i>	Riboflavin synthase alpha chain
Thioredoxin, glutaredoxin, and glutathione			
763	0824	<i>trxA</i>	Thioredoxin
764	0825	<i>trxB</i>	Thioredoxin reductase
1046	1118	<i>ggt</i>	Gamma-glutamyl transpeptidase
1091	1164	<i>trxB</i>	Thioredoxin reductase
1351	1458		Thioredoxin
Thiamine			
781	0843	<i>thiE</i>	Thiamine phosphate pyrophosphorylase
782	0844	<i>thiD</i>	Phosphomethylpyrimidine kinase
783	0845	<i>thiM</i>	Hydroxyethylthiazole kinase
Pyridine nucleotides			
312	0329	<i>nadE</i>	NH <sub>3</sub> -dependent NAD <sup>+</sup> synthetase
1273	1355	<i>nadC</i>	Nicotinate-nucleotide pyrophosphorylase
1274	1356	<i>nadA</i>	Quinolinate synthetase
Cell envelope			
Membranes and porins			
7	0009		Outer membrane protein
21	0025		Outer membrane protein

Continued on following page

TABLE 1—Continued

Gene no. in:		Gene name	Function
J99	26695		
73	0078		Outer membrane protein
	0079		
117	0127		Outer membrane protein
195	0209		Outer membrane protein
212	0227		Outer membrane protein
214	0229	<i>hopA</i>	Outer membrane protein—porin
237	0252		Outer membrane protein
238	0253, 0254		Outer membrane protein
307	0324		Outer membrane protein
342	1083		Outer membrane protein
359	1066		Outer membrane protein
424	0472		Outer membrane protein
429	0477		Outer membrane protein
438	0486		Outer membrane protein
439	0487		Outer membrane protein
456	0506		Outer membrane protein
514	0567		Inner membrane protein
581	0638		Outer membrane protein
600	0655		Outer membrane protein
614	0671		Outer membrane protein
634	0694		Outer membrane protein
645	0706	<i>hopE</i>	Outer membrane protein—porin
649	0710		Outer membrane protein
659	0722		Outer membrane protein
662	0725		Outer membrane protein
663	0726		Outer membrane protein
719	0782		Outer membrane protein
725	0788		Outer membrane protein
732	0796		Outer membrane protein
777	0839		Outer membrane protein
810	0876	<i>frpB</i>	Iron-regulated outer membrane protein
833	1243	<i>babB</i>	Outer membrane protein—adhesin
848	0912	<i>hopC</i>	Outer membrane protein—porin
849	0913	<i>hopB</i>	Outer membrane protein—porin
850	0914		Outer membrane protein
851	0915, 0916	<i>frpB</i>	Iron-regulated outer membrane protein
857	0923		Outer membrane protein
870			Outer membrane protein
1008	0373		Outer membrane protein
1022	0358		Outer membrane protein
1034	1107		Outer membrane protein
1040	1113		Outer membrane protein
1054	1125		Outer membrane protein
1083	1156		Outer membrane protein
1084	1157		Outer membrane protein
1094	1167		Outer membrane protein
1103	1177		Outer membrane function
1138	1215, 1216	<i>imp</i>	Role in outer membrane permeability
1164	0896	<i>babA</i>	Outer membrane protein—adhesin
1261	1342		Outer membrane protein
1343	1450		Inner membrane protein
1346	1453		Outer membrane protein
1349	1456	<i>lpp20</i>	Conserved lipoprotein
1360	1467		Outer membrane protein
1362	1469		Outer membrane protein
1394	1501		Outer membrane protein
1405	1512	<i>frpB</i>	Iron-regulated outer membrane protein
1432	1395		Outer membrane protein
1472	1564		Outer membrane protein
1479	1571		Outer membrane protein
Murein sacculus and peptidoglycan			
445	0493	<i>mraY</i>	Phospho- <i>N</i> -acetylmuramoyl-pentapeptide-transferase
446	0494	<i>murD</i>	UDP- <i>N</i> -acetylmuramoylalanine-D-glutamate ligase
496	0549	<i>murI</i>	Glutamate racemase
544	0597		Penicillin-binding protein
567	0623	<i>murC</i>	UDP- <i>N</i> -acetylmuramate-alanine ligase

Continued on following page



TABLE 1—Continued

Gene no. in:		Gene name	Function
J99	26695		
590	0645		Lytic murein transglycosylase
593	0648	<i>murA</i>	UDP- <i>N</i> -acetylglucosamine enolpyruvyltransferase
675	0738	<i>ddl</i>	D-Alanine-D-alanine ligase
677	0740	<i>murF</i>	D-Alanyl-D-alanine-adding enzyme
709	0772	<i>amiA</i>	Probable <i>N</i> -acetylmuramoyl-L-alanine amidase
876	0941	<i>abr</i>	Alanine racemase, biosynthetic
1082	1155	<i>murG</i>	UDP- <i>N</i> -acetylglucosamine- <i>N</i> -acetylmuramyl-(pentapeptide) pyrophosphoryl-undecaprenoln-acetylglucosaminetransferase
1313	1418	<i>murB</i>	UDP- <i>N</i> -acetylenolpyruvoyl glucosamine reductase
1387	1494	<i>murE</i>	UDP- <i>N</i> -acetylmuramyl-tripeptide synthetase
1464	1556		Penicillin-binding protein
1473	1565		Penicillin-binding protein
Surface polysaccharides, lipopolysaccharides, and antigens			
3	0003	<i>kdsA</i>	3-Deoxy-D-manno-octulosonic acid 8-phosphate synthase
37	0043	<i>manA/manC</i>	Phosphomannose isomerase/GDP-mannose pyrophosphorylase
38	0044	<i>Gmd</i>	GDP-D-mannose dehydratase
39	0045		Sugar nucleotide biosynthesis
70	0075	<i>glmM</i>	Phosphoglucosamine mutase
86	0093,0094		$\alpha$ -(1,2)-Fucosyltransferase
147	0159		Lipopolysaccharide biosynthesis protein
166	0178	<i>neuB</i>	Sialic acid synthase
182	0196	<i>lpxD</i>	UDP-3- <i>O</i> -[3-hydroxymyristoyl] glucosamine <i>N</i> -acetyltransferase
194	0208		Lipopolysaccharide biosynthesis protein
215	0230	<i>kdsB</i>	3-Deoxy-manno-octulosonate cytidyltransferase
264	0279	<i>waaC</i>	Lipopolysaccharide heptosyltransferase-1
265	0280	<i>waaM</i>	Lipid A biosynthesis acyltransferase
309	0326	<i>neuA</i>	Acylneuraminate cytidyltransferase
311	0328	<i>lpxK</i>	Tetraacyldisaccharide-1-p 4'-kinase
373	1052	<i>lpxC</i>	UDP-3- <i>O</i> -[3-hydroxymyristoyl]- <i>N</i> -acetylglucosamine deacetylase
562			Lipopolysaccharide biosynthesis protein
563	0619		Lipopolysaccharide biosynthesis protein
596	0651	<i>fucT</i>	$\alpha$ -(1,3)-Fucosyltransferase
620	0679		Lipopolysaccharide biosynthesis protein
741	0805		Lipopolysaccharide biosynthesis protein
765	0826		Lipopolysaccharide biosynthesis protein
778	0840		Sugar nucleotide biosynthesis protein
791	0857	<i>gmhA</i>	Phosphoheptose isomerase
792	0858	<i>waaE</i>	ADP-D-glycero-D-mannoheptose synthase
793	0859	<i>gmhD</i>	ADP-L-glycero-D-mannoheptose-6-epimerase
801	0867	<i>lpxB</i>	Lipid-A-disaccharide synthase
820			Lipopolysaccharide biosynthesis protein
891	0957	<i>waaA</i>	3-Deoxy-D-manno-octulosonic-acid transferase
963	0421		Polysaccharide biosynthesis protein
1002	0379	<i>fucU</i>	$\alpha$ -(1,3)-Fucosyltransferase
1015	0366		Sugar nucleotide biosynthesis
1020	0360	<i>galE</i>	UDP-glucose 4-epimerase
1031	1105		Lipopolysaccharide biosynthesis protein
1032			Lipopolysaccharide biosynthesis protein
1116	1191	<i>waaF</i>	ADP-heptose-lipopolysaccharide heptosyltransferase II
1196	1275	<i>manB</i>	Phosphomannomutase
1289	1375	<i>lpxA</i>	UDP- <i>N</i> -acetylglucosamine acyltransferase
1311	1416		Lipopolysaccharide biosynthesis protein
1368	1475	<i>kdtB</i>	Lipopolysaccharide core biosynthesis protein
1488	1581	<i>wecA</i>	Undecaprenyl-phosphate- $\alpha$ - <i>N</i> -acetylglucosaminyltransferase
Surface structures			
107	0115	<i>flaB</i>	Flagellin B
159	0173	<i>flhR</i>	Flagellar biosynthesis protein
217	0232		Motility protein
231	0246	<i>flgI</i>	Flagellar P-ring protein
280	0295	<i>flgL</i>	Flagellar hook-associated protein 3 (hap3)
308	0325	<i>flgH</i>	Flagellar L-ring protein precursor (basal body L-ring protein)

Continued on following page

TABLE 1—Continued

Gene no. in:		Gene name	Function
J99	26695		
310	0327	<i>flaG</i>	Flagellar biosynthesis protein
325	0351	<i>fliF</i>	Flagellar M-ring protein
326	0352	<i>fliG</i>	Flagellar motor switch protein
327	0353	<i>fliH</i>	Flagellar export apparatus
333	1092	<i>flgG</i>	Flagellar basal-body rod protein
383	1041	<i>flhA</i>	Flagellar biosynthesis protein
389	1035	<i>flhF</i>	Flagellar biosynthesis protein
393	1031	<i>fliM</i>	Flagellar motor switch protein
394	1030		Flagellar motor switch protein
444	0492		Paralogue of <i>hpaA</i>
531	0584	<i>fliN</i>	Flagellar motor switch protein
548	0601	<i>flaA</i>	Flagellin A
625	0684, 0685	<i>fliP</i>	Flagellar biosynthesis protein
688	0751		Flagellin protein
689	0752	<i>fliD</i>	Flagellar hook-associated protein 2 (hap2)
690	0753	<i>fliS</i>	Flagellar protein
707	0770	<i>flhB</i>	Flagellar biosynthesis protein
733	0797	<i>hpaA</i>	Neuraminylactose-binding hemagglutinin precursor
745	0809	<i>fliL</i>	Flagellar biosynthesis protein
751	0815	<i>motA</i>	Chemotaxis protein (motility protein a)
752	0816	<i>motB</i>	Flagellar motor protein
804	0870	<i>flgE</i>	Flagellar hook protein
843	0907		Flagellar biosynthesis protein
844	0908		Flagellar basal-body/rod/hook protein
971	0410		Paralogue of <i>hpaA</i>
1047	1119	<i>flgK</i>	Flagellar hook-associated protein 1 (Hap1)
1117	1192		Motility protein
1195	1274	<i>pflA</i>	Flagellar functional protein
1314	1419	<i>fliQ</i>	Flagellar biosynthesis protein
1315	1420	<i>fliI</i>	Flagellum-specific ATP synthase
1355	1462		Motility protein
1465	1557	<i>fliE</i>	Flagellar hook-basal-body complex protein
1466	1558	<i>flgC</i>	Flagellar basal-body rod protein
1467	1559	<i>flgB</i>	Flagellar basal-body rod protein
1483	1575	<i>flhB</i>	Flagellar biosynthesis protein
1492	1585	<i>flgG</i>	Flagellar basal-body rod protein (distal rod protein)
Cellular processes			
General			
4	0004	<i>icfA</i>	Carbonic anhydrase
161	0175		Peptidyl-prolyl <i>cis-trans</i> isomerase
183	0197	<i>metK</i>	S-Adenosylmethionine synthetase
228	0243	<i>napA</i>	Neutrophil-activating protein A
466	0517	<i>era</i>	GTP-binding protein
678	0741		HIT family protein
865	0930	<i>surE</i>	Stationary-phase protein
977	0404		HIT family protein
1112	1186		Carbonic anhydrase
Cell division			
314	0331	<i>minD</i>	Cell division inhibitor
315	0332	<i>minE</i>	Cell division topological specificity factor
335	1090	<i>ftsK</i>	Septum formation protein
680	0743	<i>rodA</i>	Rod shape-determining protein
912	0978	<i>ftsA</i>	Septum formation protein
913	0979	<i>ftsZ</i>	GTPase in circumferential ring formation
1086	1159	<i>fic</i>	cAMP-induced cell filamentation protein
1287	1373	<i>mreB</i>	Rod shape-determining protein
1468	1560	<i>rodA</i>	Rod shape-determining protein
Cell killing			
274	0289		Vacuolating cytotoxin (VacA) paralogue
339	1086	<i>hlyA</i>	Hemolysin
556	0609, 0610		Vacuolating cytotoxin (VacA) paralogue
819	0887	<i>vacA</i>	Vacuolating cytotoxin
856	0922		Vacuolating cytotoxin (VacA) paralog
Cag island proteins and transposable elements			
15	0017	<i>virB4a</i>	DNA transfer protein

Continued on following page

TABLE 1—Continued

Gene no. in:		Gene name	Function
J99	26695		
36	0041, 0042	<i>virB10</i>	DNA transfer protein
469	0520	<i>orf6</i>	Cag island protein
470	0521	<i>orf7</i>	Cag island protein
471	0522	<i>orf8</i>	Cag island protein
472	0523	<i>orf9</i>	Cag island protein
473	0524	<i>virD4</i>	Cag island protein, DNA transfer protein
474	0525	<i>virB11a</i>	Cag island protein, DNA transfer protein
475	0526	<i>orf12</i>	Cag island protein
476	0527	<i>orf13/14</i>	Cag island protein
477	0528	<i>orf15</i>	Cag island protein
478	0529	<i>orf16</i>	Cag island protein
479	0530	<i>orf17</i>	Cag island protein
480	0531	<i>orf18</i>	Cag island protein
481	0532	<i>cagT</i>	Cag island protein
482	0534	<i>cagS</i>	Cag island protein
483	0535	<i>cagQ</i>	Cag island protein
484	0536	<i>cagP</i>	Cag island protein
485	0537	<i>cagM</i>	Cag island protein
486	0538	<i>cagN</i>	Cag island protein
487	0539	<i>cagL</i>	Cag island protein
488	0540	<i>cagI</i>	Cag island protein
489	0541	<i>cagH</i>	Cag island protein
490	0542	<i>cagG</i>	Cag island protein
491	0543	<i>cagF</i>	Cag island protein
492	0544	<i>cagE</i>	DNA transfer protein ( <i>Agrobacterium</i> VirB4 homologue)
493	0545	<i>cagD</i>	Cag island protein
494	0546	<i>cagC</i>	Cag island protein
495	0547	<i>cagA</i>	Cag island protein, cytotoxicity-associated immunodominant antigen
826		<i>tnpB</i>	IS606 transposase
827		<i>tnpA</i>	IS606 transposase
917		<i>virB4b</i>	DNA transfer protein
918		<i>virB4c</i>	DNA transfer protein
1279	1361	<i>comEC</i>	DNA transfer protein
1316	1421	<i>virB11b</i>	DNA transfer protein
Chaperones			
8	0010	<i>groEL</i>	60-kDa chaperone
9	0011	<i>groES</i>	10-kDa chaperone
101	0109	<i>dnaK</i>	70-kDa chaperone
102	0110	<i>grpE</i>	24-kDa chaperone
196	0210	<i>hspG</i>	90-kDa chaperone
400	1024	<i>dnaJ2</i>	Cochaperone with DnaK
861	0927	<i>hspX</i>	Stress protein
1252	1332	<i>dnaJ1</i>	Cochaperone with DnaK
Detoxification			
809	0875	<i>katA</i>	Catalase
991	0390	<i>tpx</i>	Thiol peroxidase
992	0389	<i>sodF</i>	Iron-dependent superoxide dismutase
1471	1563	<i>tsaA</i>	Peroxidase
Protein and peptide secretion			
69	0074	<i>lspA</i>	Lipoprotein signal peptidase
168	0180	<i>lnt</i>	Apolipoprotein <i>N</i> -acyltransferase
329	0355	<i>lepA</i>	GTP-binding protein
523	0576	<i>lepB</i>	Signal peptidase I
700	0763	<i>ftsY</i>	Functional homolog of srp receptor
723	0786	<i>secA</i>	Preprotein translocase subunit
731	0795	<i>tig</i>	Trigger factor
889	0955	<i>lgt</i>	Prolipoprotein diacylglycerol transferase
1079	1152	<i>ffh</i>	Signal recognition particle protein
1126A	1203A	<i>secE</i>	Preprotein translocase subunit
1176	1255	<i>secG</i>	Protein export membrane protein
1220	1300	<i>secY</i>	Preprotein translocase subunit
1449	1550	<i>secD</i>	Protein export membrane protein
1450	1549	<i>secF</i>	Protein export membrane protein
Phosphorus compounds			
413	1010	<i>ppk</i>	Polyphosphate kinase
564	0620	<i>ppa</i>	Inorganic pyrophosphatase

Continued on following page



TABLE 1—Continued

Gene no. in:		Gene name	Function
J99	26695		
Polyamine biosynthesis			
18	0020	<i>nspC</i>	Carboxynorspermidine decarboxylase
771	0832	<i>speE</i>	Spermidine synthase
962	0422	<i>speA</i>	Arginine decarboxylase
Urea			
62	0067	<i>ureH</i>	Urease accessory protein
63	0068	<i>ureG</i>	Urease accessory protein
64	0069	<i>ureF</i>	Urease accessory protein
65	0070	<i>ureE</i>	Urease accessory protein
67	0072	<i>ureB</i>	Urease beta subunit
68	0073	<i>ureA</i>	Urease alpha subunit
DNA replication			
10	0012	<i>dnaG</i>	DNA primase
108	0116	<i>topA</i>	DNA topoisomerase I
199	0213	<i>gidA</i>	Glucose-inhibited division protein A
362	1063	<i>gidB</i>	Glucose-inhibited division protein B
452	0500	<i>dnaN</i>	DNA polymerase III, beta chain
453	0501	<i>gyrB</i>	DNA gyrase subunit B
558	0615	<i>lig</i>	DNA ligase
641	0701	<i>gyrA</i>	DNA gyrase subunit A
655	0717	<i>dnaX</i>	DNA polymerase III subunits gamma and tau
847	0911		ATP-dependent helicase
919		<i>topA</i>	Topoisomerase I
931		<i>topA</i>	Topoisomerase I
994	0387	<i>priA</i>	Primosomal protein n' (replication factor y)
1152	1231	<i>holB</i>	DNA polymerase III subunit delta'
1166	1245	<i>ssb</i>	Single-strand binding protein
1280	1362	<i>dnaB</i>	Replicative DNA helicase
1353	1460	<i>dnaE</i>	DNA polymerase III, alpha chain
1363	1470	<i>polA</i>	DNA polymerase I
1371	1478	<i>rep</i>	ATP-dependent DNA helicase
1412	1523	<i>recG</i>	ATP-dependent DNA helicase
1417	1529	<i>dnaA</i>	Chromosomal replication initiator protein
1438	1387		DNA polymerase III
1446	1553	<i>pcrA</i>	ATP-dependent helicase
DNA restriction, modification, recombination, and repair			
43	0050		Type II DNA modification enzyme (methyltransferase)
44			Type II DNA modification enzyme (methyltransferase)
45			Type II DNA modification enzyme (methyltransferase)
46			Type II restriction enzyme
84	0091		Type II restriction enzyme
85	0092		Type II DNA modification enzyme (methyltransferase)
130	0142	<i>mutY</i>	A/G-specific adenine glycosylase
141	0153	<i>recA</i>	Recombination protein
164			Restriction enzyme
209	0223	<i>radA</i>	DNA repair protein
243	0259	<i>xseA</i>	Exodeoxyribonuclease large subunit
244	0260		Type II DNA modification enzyme (methyltransferase)
248	0263		Type II DNA modification enzyme (methyltransferase)
306	0323		Endonuclease
322	0348	<i>recJ</i>	Single-stranded-DNA-specific exonuclease
366	1059	<i>rwvB</i>	Holliday junction DNA helicase
414		<i>hsdSI</i>	Type I restriction enzyme (specificity subunit)
415	0463	<i>hsdMI</i>	Type I restriction enzyme (modification subunit)
416	0464	<i>hsdRI</i>	Type I restriction enzyme (restriction subunit)
430	0478		Type II DNA modification enzyme (methyltransferase)
433	0481		Type II DNA modification enzyme (methyltransferase)
435	0483		Type II DNA modification enzyme (methyltransferase)
532	0585	<i>nth</i>	Endonuclease III
549	0602		Endonuclease III
565	0621	<i>mutS</i>	DNA mismatch repair protein
606	0661	<i>rnhA</i>	RNase HI
617	0675		Integrase-recombinase protein ( <i>xerCD</i> family)
618	0676	<i>ogt</i>	Methylated-DNA-protein-cysteine methyltransferase
629			Type II DNA modification enzyme (methyltransferase)

Continued on following page

TABLE 1—Continued

Gene no. in:		Gene name	Function
J99	26695		
630			Type II restriction enzyme
644	0705	<i>uvrA</i>	Excinuclease ABC subunit A
726		<i>hsdS</i>	Type I restriction enzyme (specificity subunit)
756			Type II DNA modification enzyme (methyltransferase)
760	0821	<i>uvrC</i>	Excinuclease ABC subunit C
784	0846	<i>hsdR2</i>	Type I restriction enzyme (restriction subunit)
785	0848, 0849	<i>hsdS2</i>	Type I restriction enzyme (specificity subunit)
786	0850	<i>hsdM2</i>	Type I restriction enzyme (modification subunit)
811	0877	<i>ruvC</i>	Crossover junction endodeoxyribonuclease
815	0883	<i>ruvA</i>	Holliday junction DNA helicase
846	0910		Type II DNA modification enzyme (methyltransferase)
859	0925	<i>recR</i>	Recombination protein
941	0995		Integrase/recombinase ( <i>xerCD</i> family)
951			Integrase/recombinase ( <i>xerCD</i> family)
1012	0369		Type II DNA modification enzyme (methyltransferase)
1041	1114	<i>uvrB</i>	Excinuclease ABC subunit B
1050	1121		Type II DNA modification enzyme (methyltransferase)
1131	1208	<i>M.HpyI</i>	Type II DNA modification enzyme (methyltransferase)
1149	1228	<i>mutT</i>	dGTP pyrophosphohydrolase
1243	1323	<i>mhB</i>	RNase HII
1266	1347	<i>ung</i>	Uracil-DNA glycosylase
1271	1352		Type II DNA modification enzyme (methyltransferase)
1284			Type II DNA modification enzyme (methyltransferase)
1295	1382		Endonuclease
1296		<i>mod1</i>	Type III DNA modification enzyme (methyltransferase)
1297		<i>res1</i>	Type III restriction enzyme
1364	1471		Type II restriction enzyme
1365	1472		Type II DNA modification enzyme (methyltransferase)
1409			Type II DNA modification enzyme (methyltransferase)
1410	1521	<i>res2</i>	Type III restriction enzyme
1411	1522	<i>mod2</i>	Type III DNA modification enzyme (methyltransferase)
1415	1526	<i>exoA</i>	Exodeoxyribonuclease
1422		<i>hsdS3</i>	Type I restriction enzyme (specificity subunit)
1423	1403	<i>hsdM3</i>	Type I restriction enzyme (modification subunit)
1424	1402	<i>hsdR3</i>	Type I restriction enzyme (restriction subunit)
1434	1393	<i>recN</i>	DNA repair protein
1442	1366		Type II restriction enzyme
Energy metabolism			
Amino acids and amines			
120	0132	<i>sdaB</i>	L-Serine/L-Threonine deaminase
279	0294	<i>aimE</i>	Aliphatic amidase
585			3-Hydroxyacid dehydrogenase
661	0723	<i>ansB</i>	L-Asparaginase II
1159	1238		Aliphatic amidase
1427	1399	<i>rocF</i>	Arginase
1428	1398	<i>ald</i>	L-Alanine dehydrogenase
ATP-proton motive force interconversion			
767	0828	<i>atpB</i>	ATP synthase F0, subunit a
1059	1131	<i>atpC</i>	ATP synthase F1, subunit epsilon
1060	1132	<i>atpD</i>	ATP synthase F1, subunit beta
1061	1133	<i>atpG</i>	ATP synthase F1, subunit gamma
1062	1134	<i>atpA</i>	ATP synthase F1, subunit alpha
1063	1135	<i>atpH</i>	ATP synthase F1, subunit delta
1064	1136	<i>atpF</i>	ATP synthase F0, subunit b
1065	1137	<i>atpX</i>	ATP synthase B'
1135	1212	<i>atpE</i>	ATP synthase F0, subunit c
Electron transport			
40	0047	<i>hypE</i>	Hydrogenase expression/formation protein
48	0056	<i>putA</i>	Proline/pyrroline-5-carboxylate dehydrogenase
132	0144	<i>fixN</i>	Cytochrome oxidase (CBB3-TYPE)
133	0145	<i>fixO</i>	Cytochrome oxidase (CBB3-TYPE)
134	0146	<i>fixQ</i>	Cytochrome oxidase (CBB3-TYPE)
135	0147	<i>fixP</i>	Cytochrome oxidase (CBB3-TYPE)
177	0191	<i>frdB</i>	Fumarate reductase
178	0192	<i>frdA</i>	Fumarate reductase
179	0193	<i>frdC</i>	Fumarate reductase

Continued on following page

TABLE 1—Continued

Gene no. in:		Gene name	Function
J99	26695		
250	0265	<i>ccdA</i>	Cytochrome <i>c</i> biogenesis protein
262	0277		Ferredoxin
459	0509	<i>glcD</i>	Glycolate oxidase
574	0631	<i>hyaA</i>	Hydrogenase, small subunit
575	0632	<i>hyaB</i>	Hydrogenase, large subunit
576	0633	<i>hyaC</i>	Hydrogenase, cytochrome subunit
577	0634	<i>hyaD</i>	Hydrogenase expression/formation protein
611	0666	<i>glpC</i>	Glycerol-3-phosphate dehydrogenase
803	0869	<i>hypA</i>	Hydrogenase expression/formation protein
835	0898	<i>hypD</i>	Hydrogenase expression/formation protein
836	0899	<i>hypC</i>	Hydrogenase expression/formation protein
837	0900	<i>hypB</i>	Hydrogenase expression/formation protein
878	0943	<i>dadA</i>	D-Amino acid dehydrogenase
895	0961	<i>gpsA</i>	Glycerol-3-phosphate dehydrogenase (NAD <sup>+</sup> )
974	0407		S/N-oxide reductase
1003	0378		Cytochrome <i>c</i> biogenesis protein
1035	1108	<i>porG</i>	Pyruvate ferredoxin oxidoreductase
1036	1109	<i>porD</i>	Pyruvate ferredoxin oxidoreductase
1037	1110	<i>porA</i>	Pyruvate ferredoxin oxidoreductase
1038	1111	<i>porB</i>	Pyruvate ferredoxin oxidoreductase
1088	1161	<i>fldA</i>	Flavodoxin
1090	1163	<i>fixS</i>	Component of cation transport for <i>cbb3</i> -type oxidase
1143	1222	<i>dld</i>	D-Lactate dehydrogenase
1148	1227		Periplasmic cytochrome <i>c</i> -553
1181	1260	<i>nuoA</i>	NADH oxidoreductase I
1182	1261	<i>nuoB</i>	NADH oxidoreductase I
1183	1262	<i>nuoC</i>	NADH oxidoreductase I
1184	1263	<i>nuoD</i>	NADH oxidoreductase I
1185	1264	<i>nuoE</i>	NADH oxidoreductase I
1186	1265	<i>nuoF</i>	NADH oxidoreductase I
1187	1266	<i>nuoG</i>	NADH oxidoreductase I
1188	1267	<i>nuoH</i>	NADH oxidoreductase I
1189	1268	<i>nuoI</i>	NADH oxidoreductase I
1190	1269	<i>nuoJ</i>	NADH oxidoreductase I
1191	1270	<i>nuoK</i>	NADH oxidoreductase I
1192	1271	<i>nuoL</i>	NADH oxidoreductase I
1193	1272	<i>nuoM</i>	NADH oxidoreductase I
1194	1273	<i>nuoN</i>	NADH oxidoreductase I
1354	1461		Cytochrome <i>c</i> peroxidase
1401	1508	<i>fixG</i>	Component of cation transport for <i>cbb3</i> -type oxidase
1459	1540	<i>petA</i>	Ubiquinol cytochrome <i>c</i> oxidoreductase, 2Fe-2S subunit
1460	1539	<i>petB</i>	Ubiquinol cytochrome <i>c</i> oxidoreductase, cytochrome <i>b</i> subunit
1461	1538	<i>petC</i>	Ubiquinol cytochrome <i>c</i> oxidoreductase, cytochrome <i>cI</i> subunit
Entner-Doudoroff pathway			
1025	1099	<i>eda</i>	2-Keto-3-deoxy-6-phosphogluconate aldolase
1026	1100	<i>edd</i>	Phosphogluconate dehydratase
Fermentation			
840	0903	<i>ackA</i>	Acetate kinase
841	0904, 0905	<i>pta</i>	Phosphotransacetylase
1030	1104		Zinc-dependent alcohol dehydrogenase
1429			Zinc-dependent alcohol dehydrogenase
Gluconeogenesis			
111	0121	<i>ppsA</i>	Phosphoenolpyruvate synthase
142	0154	<i>eno</i>	Enolase
162	0176	<i>fba</i>	Fructose-bisphosphate aldolase
180	0194	<i>tpi</i>	Triose-phosphate isomerase
855	0921	<i>gap</i>	Glyceraldehyde-3-phosphate dehydrogenase
908	0974	<i>pgm</i>	2,3-Bisphosphoglycerate-independent phosphoglycerate mutase
1093	1166	<i>pgi</i>	Glucose-6-phosphate isomerase
1264	1345	<i>pgk</i>	Phosphoglycerate kinase
1265	1346	<i>gap</i>	Glyceraldehyde 3-phosphate dehydrogenase
1440	1385	<i>fbp</i>	Fructose-1,6-bisphosphatase
Phosphopentose pathway			
337	1088	<i>tktA</i>	Transketolase
521	0574	<i>rpi</i>	Ribose 5-phosphate isomerase

Continued on following page

TABLE 1—Continued

Gene no. in:		Gene name	Function
J99	26695		
1027	1101	<i>zwf</i>	Glucose-6-phosphate-1-dehydrogenase
1388	1495	<i>tal</i>	Transaldolase
1439	1386	<i>rpe</i>	Ribulose-phosphate-3-epimerase
Sugars			
1029	1103	<i>glk</i>	Glucokinase
Tricarboxylic acid cycle			
22	0026	<i>glcA</i>	Citrate synthase
23	0027	<i>icd</i>	Isocitrate dehydrogenase
536	0588	<i>oorD</i>	Subunit of 2-oxoglutarate oxidoreductase
537	0589	<i>oorA</i>	Subunit of 2-oxoglutarate oxidoreductase
538	0590	<i>oorB</i>	Subunit of 2-oxoglutarate oxidoreductase
539	0591	<i>oorC</i>	Subunit of 2-oxoglutarate oxidoreductase
716	0779	<i>acnB</i>	Aconitate hydratase
1245	1325	<i>fumC</i>	Fumarase
Other			
88	0096		Keto-acid dehydrogenase
586	0642		Oxidoreductase
888	0954		Aldehyde dehydrogenase
1023	0357		Oxidoreductase
1028	1102		Dehydrogenase
1345	1452	<i>thdF</i>	Thiophene/furan oxidation protein
Fatty acid and phospholipid metabolism			
83	0090	<i>fabD</i>	Malonyl-CoA-ACP transacylase
176	0190		Cardiolipin synthase
181	0195	<i>fabI</i>	Enoyl-ACP reductase
187	0201	<i>plsX</i>	Fatty acid/phospholipid synthesis protein
188	0202	<i>fabH</i>	β-Ketoacyl-ACP synthase III
201	0215	<i>cdsA</i>	CDP-diacylglycerol synthase
354	1071	<i>pssA</i>	Phosphatidylserine synthase
407	1016	<i>pgsA</i>	Phosphatidylglycerophosphate synthase
409	1014		Short-chain dehydrogenase
451	0499	<i>pldA</i>	Phospholipase A <sub>1</sub>
504	0557	<i>accA</i>	Acetyl-CoA carboxylase subunit A
505	0558	<i>fabB</i>	β-Ketoacyl-ACP synthase I
506	0559	<i>acpP</i>	ACP
508	0561	<i>fabG</i>	Acetyl-CoA carboxylase subunit A
636	0692	<i>scoB</i>	3-Oxoacid CoA-transferase, subunit B
637	0691	<i>scoA</i>	3-Oxoacid CoA-transferase, subunit A
638	0690	<i>thl</i>	Acetyl-CoA acetyltransferase
640	0700	<i>dgkA</i>	Diacylglycerol kinase
674	0737	<i>pgpA</i>	Phosphatidylglycerophosphatase A
744	0808	<i>acpS</i>	Holo-ACP synthase
805	0871	<i>cdh</i>	CDP-diacylglycerol pyrophosphatase
884	0950	<i>accD</i>	Acetyl-CoA carboxylase subunit B
968	0416	<i>cfa</i>	Cyclopropane fatty acid synthase
1010	0371	<i>accB</i>	Biotin carboxyl carrier protein
1011	0370	<i>accC</i>	Biotin carboxylase
1267	1348	<i>plsC</i>	1-Acyl-SN-glycerol-3-phosphate acyltransferase
1275	1357	<i>psd</i>	Phosphatidylserine decarboxylase
1290	1376	<i>fabZ</i>	Hydroxymyristoyl-ACP dehydratase
Purines, pyrimidines, nucleosides and nucleotides			
2'-Deoxyribonucleotide metabolism			
621	0680	<i>nrdA</i>	Ribonucleoside-diphosphate reductase 1 alpha chain
799	0865	<i>dut</i>	Deoxyuridine 5'-triphosphate nucleotidohydrolase
1009	0372	<i>dcd</i>	Deoxycytidine triphosphate deaminase
1016	0364	<i>nrdB</i>	Ribonucleoside-diphosphate reductase 1 beta chain
Purine ribonucleotide biosynthesis			
1039	1112	<i>purB</i>	Adenylosuccinate lyase
1140	1218	<i>purD</i>	Glycinamide ribonucleotide synthetase
1327	1434	<i>purU</i>	Formyltetrahydrofolate hydrolase
Pyrimidine ribonucleotide biosynthesis			
5	0005	<i>pyrF</i>	Orotidine 5'-phosphate decarboxylase
184	0198	<i>ndk</i>	Nucleoside diphosphate kinase
251	0266	<i>pyrC1</i>	Dihydroorotase
323	0349	<i>pyrG</i>	CTP synthase

Continued on following page

TABLE 1—Continued

Gene no. in:		Gene name	Function
J99	26695		
341	1084	<i>pyrB</i>	Aspartate carbamoyltransferase catalytic chain
412	1011	<i>pyrD</i>	Dihydroorotate dehydrogenase
528	0581	<i>pyrC2</i>	Dihydroorotase
714	0777	<i>pyrH</i>	Uridylate kinase
853	0919	<i>pyrA1</i>	Carbamoyl-phosphate synthase large chain
1158	1237	<i>pyrA2</i>	Carbamoyl-phosphate synthase small chain
1178	1257	<i>pyrE</i>	Orotate phosphoribosyltransferase
Salvage and interconversion of nucleosides and nucleotides			
96	0104	<i>cpdB</i>	2',3'-Cyclic nucleotide 2'-phosphodiesterase
239	0255	<i>purA</i>	Adenylosuccinate synthetase
304	0321	<i>gmk</i>	Guanylate kinase
519	0572	<i>apt</i>	Adenine phosphoribosyltransferase
561	0618	<i>adk</i>	Adenylate kinase
672	0735	<i>gpt</i>	Xanthine-guanine phosphoribosyltransferase
679	0742	<i>prsA</i>	Phosphoribosyl pyrophosphate synthetase
768	0829	<i>guaB</i>	Inosine-5'-monophosphate dehydrogenase
790	0854	<i>guaC</i>	GMP reductase
972	0409	<i>guaA</i>	GMP synthetase
1104	1178	<i>deoD</i>	Purine nucleoside phosphorylase
1105	1179	<i>deoB</i>	Phosphopentomutase
1367	1474	<i>tmk</i>	Thymidylate kinase
Sugar-nucleotide biosynthesis and conversions			
591	0646	<i>galU</i>	UTP-glucose-1-phosphate uridylyltransferase
624	0683	<i>glmU</i>	UDP-N-acetylglucosamine pyrophosphorylase
1420	1532	<i>glmS</i>	Glutamine fructose-6-phosphate aminotransferase
Regulatory functions			
General			
41	0048		Transcriptional regulator
81	0088	<i>rpoD</i>	RNA polymerase sigma 70 factor
151	0164, 0165		Histidine kinase sensor protein
152	0166		Transcriptional regulator
229	0244		Histidine kinase sensor protein
263	0278	<i>gppA</i>	Guanosine-5'-triphosphate,3'-diphosphate pyrophosphatase
381	1043		Transcriptional regulator
392	1032	<i>flaA</i>	RNA polymerase sigma 28 factor
397	1027	<i>fur</i>	Ferric uptake regulation protein
399	1025		Transcriptional regulator
403	1021		Transcriptional regulator
643	0703		Transcriptional regulator
652	0714	<i>rpoN</i>	RNA polymerase sigma-54 factor
664	0727		Transcriptional regulator
712	0775	<i>spoT</i>	Guanosine-3',5'-bis(diphosphate)-3'-pyrophosphohydrolase
981	0400	<i>lytB</i>	Lysis tolerance protein
1207	1287		Transcriptional regulator
1282	1364		Histidine kinase sensor protein
1283	1365		Transcriptional regulator
1335	1442	<i>csrA</i>	Carbon storage regulator
1443	1365		Transcriptional regulator
1480	1572	<i>dniR</i>	Regulatory protein
Chemotaxis and motility			
17	0019	<i>cheV1</i>	Chemotaxis protein
75	0082		MCP
91	0099		MCP
95	0103		MCP
358	1067	<i>cheY</i>	Response regulator
546	0599		MCP
559	0616	<i>cheV2</i>	Chemotaxis protein
988	0393	<i>cheV3</i>	Chemotaxis protein
989	0392	<i>cheA</i>	Histidine kinase
990	0391	<i>cheW</i>	Histidine kinase-MCP coupling protein
Transcription			
Degradation of RNA			
1136	1213	<i>pnp</i>	Polyribonucleotide nucleotidyltransferase
1169	1248	<i>vacB</i>	RNase II family protein

Continued on following page



TABLE 1—Continued

Gene no. in:		Gene name	Function
J99	26695		
1299	1407	<i>rhn</i>	RNase N
DNA-dependent RNA polymerase			
1121	1198	<i>rpoB</i>	DNA-directed RNA polymerase, beta subunit
1213	1293	<i>rpoA</i>	DNA-directed RNA polymerase, alpha subunit
1458	1541	<i>mfd</i>	Transcription-repair coupling factor
Transcription factors			
1	0001	<i>nusB</i>	Transcription termination
497	0550	<i>rho</i>	Transcription termination factor
800	0866	<i>greA</i>	Transcription elongation factor (transcript cleavage factor)
1126	1203	<i>nusG</i>	Transcription antitermination protein
1407	1514	<i>nusA</i>	N utilization substance protein A
RNA processing			
583	0640	<i>pcnB</i>	Polynucleotide adenylyltransferase
607	0662	<i>rnc</i>	RNase III
Translation			
Aminoacyl-tRNA synthetases			
113	0123	<i>thrS</i>	Threonyl-tRNA synthetase
170	0182	<i>lysS</i>	Lysyl-tRNA synthetase
223	0238	<i>proS</i>	Prolyl-tRNA synthetase
302	0319	<i>argS</i>	Arginyl-tRNA synthetase
428	0476	<i>gliX</i>	Glutamyl-tRNA synthetase
560	0617	<i>aspS</i>	Aspartyl-tRNA synthetase
588	0643	<i>gliX</i>	Glutamyl-tRNA synthetase
711	0774	<i>tyrS</i>	Tyrosyl-tRNA synthetase
818	0886	<i>cysS</i>	Cysteinyl-tRNA synthetase
894	0960	<i>glyQ</i>	Glycyl-tRNA synthetase alpha chain
906	0972	<i>glyS</i>	Glycyl-tRNA synthetase beta chain
967	0417	<i>metG</i>	Methionyl-tRNA synthetase
978	0403	<i>pheS</i>	Phenylalanyl-tRNA synthetase alpha chain
979	0402	<i>pheT</i>	Phenylalanyl-tRNA synthetase beta chain
1080	1153	<i>valS</i>	Valyl-tRNA synthetase
1115	1190	<i>hisS</i>	Histidyl-tRNA synthetase
1162	1241	<i>alaS</i>	Alanyl-tRNA synthetase
1174	1253	<i>trpS</i>	Tryptophanyl-tRNA synthetase
1317	1422	<i>ileS</i>	Isoleucyl-tRNA synthetase
1373	1480	<i>serS</i>	Seryl-tRNA synthetase
1452	1547	<i>leuS</i>	Leucyl-tRNA synthetase
Degradation of proteins, peptides and glycopeptides			
29	0033	<i>clpA</i>	ATP-dependent protease, ATP-binding subunit
155	0169		Protease
249	0264	<i>clpB</i>	Heat shock protein
271	0286	<i>ftsH</i>	ATP-dependent Zn metalloproteinase
356	1069	<i>ftsH</i>	ATP-dependent Zn metalloproteinase
387	1037	<i>pepQ</i>	Proline peptidase
405	1019	<i>htrA</i>	Protease DO
411	1012		Zn protease
422	0470	<i>pepF</i>	Oligopeptidase
464	0515	<i>hslV</i>	Heat shock protein
465	0516	<i>hslU</i>	Heat shock protein
517	0570	<i>pepA</i>	Aminopeptidase
602	0657		Processing protease
603	0658	<i>gatB</i>	Glu-tRNA amidotransferase, subunit B
730	0794	<i>clpP</i>	ATP-dependent protease, proteolytic subunit
769	0830	<i>gatA</i>	Glu-tRNA amidotransferase, subunit A
909	0975	<i>gatC</i>	Glu-tRNA amidotransferase, subunit C
999	0382		Zn-metallo protease
1269	1350	<i>prc</i>	Carboxyl-terminal protease
1288	1374	<i>clpX</i>	ATP-dependent protease, ATP-binding subunit
1293	1379	<i>lon</i>	ATP-dependent protease 1a
1328	1435	<i>sppA</i>	Protease
1491	1584	<i>ydiE</i>	O-Sialoglycoprotein endopeptidase
Nucleoproteins			
774	0835		DNA-binding protein HU
1052	1123	<i>styD</i>	FKBP-type peptidyl-prolyl <i>cis-trans</i> isomerase

Continued on following page

TABLE 1—Continued

Gene no. in:		Gene name	Function
J99	26695		
Protein modification			
210	0224		Peptide methionine sulfoxide reductase
729	0793	<i>def</i>	Polypeptide deformylase
1017	0363	<i>pcm</i>	Protein-L-isoaspartate <i>O</i> -methyltransferase
1219	1299	<i>map</i>	Methionine aminopeptidase
1334	1441	<i>ppiA</i>	Peptidyl-prolyl <i>cis-trans</i> isomerase
Ribosomal proteins, synthesis and modification			
71	0076	<i>rpsT</i>	30S ribosomal protein S20
76	0083	<i>rpsI</i>	30S ribosomal protein S9
77	0084	<i>rplM</i>	50S ribosomal protein L13
115	0125	<i>rpmI</i>	50S ribosomal protein L35
116	0126	<i>rplT</i>	50S ribosomal protein L20
186	0200	<i>rpmF</i>	50S ribosomal protein L32
281	0296	<i>rplU</i>	50S ribosomal protein L21
282	0297	<i>rpmA</i>	50S ribosomal protein L27
357	1068	<i>prmA</i>	Ribosomal protein L11 methyltransferase
378	1047	<i>rbfA</i>	Ribosome-binding factor A
384	1040	<i>rpsO</i>	30S ribosomal protein S15
443	0491	<i>rpmB</i>	50S ribosomal protein L28
463	0514	<i>rplI</i>	50S ribosomal protein L9
498	0551	<i>rpmE</i>	50S ribosomal protein L31
509	0562	<i>rpsU</i>	30S ribosomal protein S21
982	0399	<i>rpsA</i>	30S ribosomal protein S1
1074	1147	<i>rplS</i>	50S ribosomal protein L19
1078	1151	<i>rpsP</i>	30S ribosomal protein S16
1119	1196	<i>rpsG</i>	30S ribosomal protein S7
1120	1197	<i>rpsL</i>	30S ribosomal protein S12
1122	1199	<i>rplL</i>	50S ribosomal protein L7/L12
1123	1200	<i>rplJ</i>	50S ribosomal protein L10
1124	1201	<i>rplA</i>	50S ribosomal protein L1
1125	1202	<i>rplK</i>	50S ribosomal protein L11
1127	1204	<i>rpmG</i>	50S ribosomal protein L33
1165	1244	<i>rpsR</i>	30S ribosomal protein S18
1167	1246	<i>rpsF</i>	30S ribosomal protein S6
1212	1292	<i>rplQ</i>	50S ribosomal protein L17
1214	1294	<i>rpsD</i>	30S ribosomal protein S4
1215	1295	<i>rpsK</i>	30S ribosomal protein S11
1217	1296	<i>rpsM</i>	30S ribosomal protein S13
1217	1297	<i>rpmJ</i>	50S ribosomal protein L36
1221	1301	<i>rplO</i>	50S ribosomal protein L15
1222	1302	<i>rpsE</i>	30S ribosomal protein S5
1223	1303	<i>rplR</i>	50S ribosomal protein L18
1224	1304	<i>rplF</i>	50S ribosomal protein L6
1225	1305	<i>rpsH</i>	30S ribosomal protein S8
1226	1306	<i>rpsN</i>	30S ribosomal protein S14
1227	1307	<i>rplE</i>	50S ribosomal protein L5
1228	1308	<i>rplX</i>	50S ribosomal protein L24
1229	1309	<i>rplN</i>	50S ribosomal protein L14
1230	1310	<i>rpsQ</i>	30S ribosomal protein S17
1231	1311	<i>rpmC</i>	50S ribosomal protein L29
1232	1312	<i>rplP</i>	50S ribosomal protein L16
1233	1313	<i>rpsC</i>	30S ribosomal protein S3
1234	1314	<i>rplV</i>	50S ribosomal protein L22
1235	1315	<i>rpsS</i>	30S ribosomal protein S19
1236	1316	<i>rplB</i>	50S ribosomal protein L2
1237	1317	<i>rplW</i>	50S ribosomal protein L23
1238	1318	<i>rplD</i>	50S ribosomal protein L4
1239	1319	<i>rplC</i>	50S ribosomal protein L3
1240	1320	<i>rpsJ</i>	30S ribosomal protein S10
1340	1447	<i>rpmH</i>	50S ribosomal protein L34
1389	1496	<i>rplY</i>	50S ribosomal protein L25
1445	1554	<i>rpsB</i>	30S ribosomal protein S2
tRNA modification			
266	0281	<i>tgt</i>	Queuine-tRNA-ribosyltransferase
363	1062	<i>queA</i>	S-Adenosylmethionine tRNA ribosyltransferase-isomerase

Continued on following page

TABLE 1—Continued

Gene no. in:		Gene name	Function
J99	26695		
1019	0361	<i>truA</i>	Pseudouridylate synthase I
1069	1141	<i>fnt</i>	Methionyl-tRNA formyltransferase
1075	1148	<i>trmD</i>	tRNA (guanine-n1)-methyltransferase
1254	1335	<i>trmU</i>	tRNA(5-methylaminomethyl-2-thiouridylate)-methyltransferase
1310	1415	<i>miaA</i>	tRNA delta(2)-isopentenylpyrophosphate transferase
1341	1448	<i>rnpA</i>	RNase protein component
1390	1497	<i>pth</i>	Peptidyl-tRNA hydrolase
1406	1513	<i>selA</i>	L-Seryl-tRNA selenium transferase
Translation factors			
72	0077	<i>prfA</i>	Peptide chain release factor 1
114	0124	<i>infC</i>	Translation initiation factor IF-3
157	0171	<i>prfB</i>	Peptide chain release factor 2 (RF-2)
163	0177	<i>efp</i>	Elongation factor P (EF-P)
232	0247	<i>deaD</i>	ATP-dependent RNA helicase dead
377	1048	<i>infB</i>	Translation initiation factor IF-2
1118	1195	<i>fusA</i>	Elongation factor G (EF-G)
1128	1205	<i>tufA</i>	Elongation factor TU (EF-TU)
1177	1256	<i>frr</i>	Ribosome recycling factor (ribosome-releasing factor [RRF])
1218	1298	<i>infA</i>	Translation initiation factor IF-1
1322	1431	<i>ksgA</i>	Dimethyladenosine transferase
1444	1555	<i>tfs</i>	Elongation factor TS (EF-TS)
Transport and binding proteins			
General			
66	0071	<i>ureI</i>	Urea transporter
167	0179		ABC transporter, ATP-binding protein
200	0214		Transporter
235	0250		ABC transporter, ATP-binding protein
236	0251		ABC transporter, permease
300	0613		ABC transporter, ATP-binding protein
343	1082	<i>msbA</i>	Multidrug resistance protein
449	0497		Transporter
450	0498		Transporter
547	0600		Secretion/efflux ABC transporter, ATP-binding protein
553	0606		Efflux transporter
554	0607		Efflux transporter
653	0715		ABC transporter, ATP-binding protein
685	0748		ABC transporter, ATP-binding protein
754	0818		Osmoprotection binding protein
757	0818		Osmoprotection binding protein
758	0819		Osmoprotection ATP-binding protein
789	0853		ABC transporter, ATP-binding protein
806	0872	<i>phnA</i>	Alkylphosphonate uptake protein
871	0936	<i>proP</i>	Proline/betaine transporter
1055	1126	<i>tolB</i>	<i>tonB</i> -independent protein-uptake protein
1057	1129	<i>exbD1</i>	Biopolymer transport protein
1058	1130	<i>exbB1</i>	Biopolymer transport protein
1107	1181		Transporter
1129	1206		ABC transporter, ATP-binding protein
1141	1220		ABC transporter, ATP-binding protein
1320	1427		Histidine-rich metal-binding protein
1321	1432		Histidine- and glutamine-rich metal-binding protein
1338	1445	<i>exbB3</i>	Biopolymer transport protein
1339	1446	<i>exbD3</i>	Biopolymer transport protein
1484	1576		ABC transporter, ATP-binding protein
1485	1577		ABC transporter, permease
Amino acids, peptides, and amines			
47	0055	<i>putP</i>	Sodium/proline symporter
121	0133	<i>sdaC</i>	L-Serine transporter
283	0298	<i>dppA</i>	Periplasmic dipeptide transport substrate-binding protein
284	0299	<i>dppB</i>	Dipeptide transport system permease protein
285	0300	<i>dppC</i>	Dipeptide transport system permease protein
286	0301	<i>dppD</i>	Dipeptide transport system ATP-binding protein
287	0302	<i>dppF</i>	Dipeptide transport system ATP-binding protein
406	1017		Amino acid permease
874	0939		Amino acid ABC transporter, permease protein

Continued on following page

TABLE 1—Continued

Gene no. in:		Gene name	Function
J99	26695		
875	0940		Amino acid ABC transporter, binding protein precursor
877	0942		Sodium/alanine symporter
1096	1169		Amino acid ABC transporter, permease protein
1097	1170		Amino acid ABC transporter, permease protein
1098	1171		Amino acid ABC transporter, ATP-binding protein
1099	1172		Amino acid ABC transporter, binding protein precursor
1172	1251		Peptide ABC transporter, ATP-binding protein
1358	1465		Amino acid ABC transporter, ATP-binding protein
1399	1506	<i>gltS</i>	Sodium/glutamate symporter
Anions			
425	0473	<i>modA</i>	Molybdenum ABC transporter, periplasmic binding protein
426	0474	<i>modB</i>	Molybdenum ABC transporter, permease
427	0475	<i>modC</i>	Molybdenum ABC transporter, ATP-binding protein
1384	1491		Phosphate permease
Carbohydrates, organic alcohols, and acids			
128	0140	<i>lldP</i>	L-Lactate permease
129	0141	<i>lldP</i>	L-Lactate permease
334	1091	<i>kgfP</i>	$\alpha$ -Ketoglutarate permease
635	0693	<i>atoE</i>	Short-chain fatty acids transporter
660	0724	<i>dcuA</i>	Anaerobic C <sub>4</sub> -dicarboxylate membrane transporter
1101	1174	<i>gluP</i>	Glucose/galactose transporter
Cations			
124	0136	<i>bcp</i>	Bacterioferritin comigratory protein
348	1077	<i>nixA</i>	High-affinity nickel transport protein
352	1073	<i>copP</i>	Copper-associated protein
353	1072	<i>copA</i>	Copper-transporting P-type ATPase
423	0471	<i>kefB</i>	Glutathione-regulated potassium efflux system protein
442	0490		Putative potassium channel protein
529	0582	<i>tonB1</i>	Siderophore-mediated iron transport protein
598	0653	<i>pfr</i>	Nonheme iron-containing ferritin
626	0686	<i>fecA1</i>	Iron(III) dicitrate transport protein
627	0687	<i>feoB</i>	Ferrous iron transport protein B
727	0791	<i>hmcT</i>	Heavy-metal cation-transporting P-type ATPase
743	0807	<i>fecA2</i>	Iron(III) dicitrate transport protein
821	0888	<i>fecE</i>	Iron(III) dicitrate transport system ATP-binding protein
822	0889	<i>fecD</i>	Iron(III) dicitrate transport system permease protein
903	0969	<i>czcA1</i>	Cation efflux system protein
904	0970	<i>czcB1</i>	Cation efflux system protein
1109	1183		Na <sup>+</sup> /H <sup>+</sup> antiporter
1248	1328	<i>czcB2</i>	Cation efflux system protein
1249	1329	<i>czcA2</i>	Cation efflux system protein
1258	1339	<i>exbB2</i>	Biopolymer transport protein
1259	1340	<i>exbD2</i>	Biopolymer transport protein
1260	1341	<i>tonB2</i>	Siderophore-mediated iron transport protein
1263	1344	<i>corA</i>	Magnesium and cobalt transport protein
1396	1503	<i>fixI</i>	Component of cation transport for <i>cbb3</i> -type oxidase
1426	1400	<i>fecA3</i>	Iron(III) dicitrate transport protein
1447	1552	<i>nhaA</i>	Na <sup>+</sup> /H <sup>+</sup> antiporter I
Nucleosides, purines, and pyrimidines			
1106	1180		Nucleoside transporter
1210	1290	<i>pnuC</i>	Nicotinamide mononucleotide transporter

<sup>a</sup> Gene numbers correspond to those in Fig. 1.

phenylalanine, methionine, and the branched-chain amino acids necessitates specific transport of these amino acids. No such specific transport systems were identified, but they may be encoded by any of the transport systems with unassigned ligand specificity.

In addition to transporting amino acids, *H. pylori* may have the ability to transport the abundant peptides which are found in the stomach. Homologues to a dipeptide transport system are present and its five genes (JHP283–287/HP0298–0302) are arranged contiguously, similar to the organization of the *dpp* operon in *E. coli* (1). There is also a single gene (JHP1172/

HP1251) which displays significant sequence similarity to an oligopeptide transporter.

The genomic sequence provides little information on the composition of polyamines in *H. pylori*, which are needed for optimal growth in most cells. The homologue of SpeA allows the conversion of arginine to agmatine in *H. pylori*. Although *H. pylori* has a homologue to *speE*, which encodes spermidine synthetase, it is unlikely that this enzyme can catalyze spermidine biosynthesis since no homologues for the enzymes that provide precursors for SpeE (SpeB to SpeD) were detected. However, *H. pylori* may be able to synthesize spermidine

TABLE 2. Annotation and classification of genes from *H. pylori* J99 and 26695

Annotation category	No. of genes in:		
	<i>H. pylori</i> J99	<i>H. pylori</i> 26695	Both strains <sup>a</sup>
Functionally classified	877	898	
Conserved with no known function	275	290	
<i>H. pylori</i> specific	343	364	
Total	1,495	1,552	
Amino acid biosynthesis	44	44	44
Biosynthesis of cofactors, etc.	60 <sup>b</sup>	59	59
Cell envelope	160	164	156 <sup>c,d,e</sup>
Cellular processes	96	113	92 <sup>c,d,e</sup>
DNA replication	23	23	21 <sup>c,e</sup>
DNA restriction-modification, etc.	66	68	51 <sup>c,d,e</sup>
Energy metabolism	104	104	102 <sup>c,d,e</sup>
Fatty acid and phospholipid metabolism	28	29	28 <sup>e</sup>
Purine and pyrimidine biosynthesis	34	34	34
Regulatory functions	32	32	31 <sup>d,f</sup>
Transcription	13	13	13
Translation	128	128	128
Transport and binding proteins	88 <sup>g</sup>	87	87
Conserved with no known function	275	290	267 <sup>c,d,e</sup>
<i>H. pylori</i> specific	343	364	288 <sup>c,d,e</sup>
Total	1,495	1,552	1,401 <sup>h</sup>

<sup>a</sup> Using J99 genes as the basis for counting.

<sup>b</sup> Includes the partial duplication of *folE* (JHP862).

<sup>c</sup> Categories which include *H. pylori* J99-specific genes (see the text for details).

<sup>d</sup> These numbers include the "split" genes based on the *H. pylori* J99 definition. There are 6 J99 genes which constitute 12 26695 genes in the cell envelope; 2 J99 genes which constitute 4 26695 genes in cellular processes; 1 J99 gene which constitutes 2 26695 genes in each of DNA restriction-modification, energy metabolism, and Regulatory functions; 5 J99 genes which constitute 7 26695 genes in conserved with no known function; and 24 J99 genes which constitute 33 26695 genes in *H. pylori* specific.

<sup>e</sup> Categories which include *H. pylori* 26695-specific genes.

<sup>f</sup> Does not include the duplication of the response regulator (JHP1283 and JHP1443).

<sup>g</sup> Includes the partial duplication of *proX* (JHP754).

<sup>h</sup> The remaining 94 genes represent the 89 J99-specific genes, *mpA/B* from IS606, and the partial or complete duplications of three genes.

through *nspC*. The product of this gene can synthesize spermidine by decarboxylating carboxyspermidine (117). It is also possible that *H. pylori* uses *nspC* for the synthesis of norspermidine, a polyamine found in *Vibrio alginolyticus*.

### Cofactors and Vitamins

Both sequenced strains of *H. pylori* have all the identified genes needed for the biosynthesis of biotin, folate, heme, molybdopterin, pantothenate, pyridoxal phosphate, riboflavin, and thioredoxin (Table 4). *H. pylori* has all the genes necessary for the synthesis of NAD with the exception of *nadB*, which encodes the aspartate oxidase subunit of quinolate synthetase in *E. coli* (138). This polypeptide is the oxygen-utilizing subunit of an enzyme which converts L-aspartate to iminoaspartate.

The absence of *NadB* is not unexpected since the mechanism by which anaerobic or microaerophilic bacteria, such as *H. pylori*, synthesize iminoaspartate is unknown and is not likely to be oxygen dependent. In addition, nicotinamide mononucleotide transporter (JHP1210/HP1290) was identified. No homologues to enzymes involved in vitamin B<sub>12</sub> and coenzyme A biosynthesis were identified. Vitamin B<sub>12</sub> is an important cofactor for certain enzymes involved in anaerobic metabolism such as methionine synthase. *H. pylori* may not need to synthesize this vitamin, since homologues to B<sub>12</sub>-requiring enzymes were not identified. Bacteria synthesize coenzyme A de novo from pantothenate. However, no homologues to known enzymes involved in its biosynthesis were identified in *H. pylori*, making this pathway unique with respect to those previously reported. The pathway for thiamine biosynthesis has not been completely defined. Some of the genes believed to be involved in thiamine synthesis were found, suggesting that *H. pylori* can make this vitamin. However, it has been reported that *H. pylori* requires thiamine for growth (118).

*H. pylori* has homologues to all of the genes necessary to produce riboflavin. A single gene in *H. pylori*, homologous to both *ribB* and *ribA* (JHP740/HP0804), encodes a bifunctional enzyme with both GTP cyclohydrolase II and 3,4-dihydroxy-2-butanone 4-phosphate synthase activities (158). In addition, *H. pylori* has a separate GTP cyclohydrolase II (RibA) homologue (JHP738/HP802) downstream from the bifunctional RibAB. Worst et al. (158) have shown that expression of RibAB but not RibA is regulated by iron limitation in *H. pylori*. The significance of this enzymatic duplication and differential gene regulation is unknown.

### Purine and Pyrimidine Biosynthesis, Salvage, and Interconversion

Enzymes for the de novo biosynthesis of purines are largely absent in both sequenced strains of *H. pylori*, which implies that this bacterium cannot synthesize purine nucleotides from formate, glycine, or serine. Genes that encode homologues for all of the purine salvage and interconversion enzymes are present (Fig. 3A). No homologues to a purine transporter were identified, although there is biochemical evidence for the transport of purine bases in *H. pylori* (107, 118). Based on similarity, it is likely that the putative transporter (JHP1106/HP1180) is specific for nucleosides, which would allow *H. pylori* to obtain purines via the salvage and interconversion pathways.

*H. pylori* possesses homologues to all of the genes necessary for the de novo synthesis of UTP and CTP, consistent with experimental results that show that radiolabeled pyrimidine nucleotide precursors are incorporated into DNA (108). Unlike *E. coli*, *H. pylori* possesses a second *pyrC* homologue encoding dihydroorotase, raising the possibility that the *H. pylori* enzyme exists as a heterodimer rather than as a homodimer as reported in *E. coli*. One of the two homologues (JHP528/HP0581) is more closely related to a *PyrC* in gram-negative organisms, whereas the other homologue (JHP251/HP0266) is more closely related to a *PyrC* in gram-positive organisms. Furthermore, *H. pylori* lacks a homologue for the regulatory

FIG. 1. Linear representation of the *H. pylori* J99 chromosome, illustrating the location of each predicted protein-coding region, rRNA gene, tRNA gene, IS605 or IS606 element and related fragment, and *NotI* endonuclease site. The predicted protein-coding regions are color coded based on functional classification (see the bottom of the figure for the code), with the direction of transcription indicated by an arrowhead. *H. pylori* J99 ORFs are numbered sequentially in red, and the corresponding homologous gene, if it exists in strain 26695, is numbered in black. The positions of the *NotI* endonuclease sites in J99 are indicated with the number of conserved nucleotides in the recognition sequence (x/8) at the corresponding position in strain 26695. The numbers associated with the tRNA symbols (inverted triangles) represent the number of tRNA genes at a specific locus. Vertical hash marks, below the linear chromosome, are located every 20 kb.



TABLE 3. Nucleotide and amino acid identity between genes common to *H. pylori* J99 and 26695

% Identity	No. (%) of predicted ORFs <sup>a</sup>	
	Nucleotide	Amino acid
100	0 (0)	41 (2.9)
98.0–99.9	8 (0.6)	269 (19.3)
96.0–97.9	249 (17.8)	359 (25.7)
94.0–95.9	566 (40.5)	279 (20.0)
92.0–93.9	306 (21.9)	169 (12.1)
90.0–91.9	89 (6.4)	86 (6.2)
85.0–89.9	81 (5.8)	77 (5.5)
<85	97 (7.0)	116 (8.3)
Total	1,396 (100)	1,396 (100)

<sup>a</sup> Genes that appear “split” by putative frameshifts in either strain have been classed as the larger ORF in the above analysis.

chain of the aspartate transcarbamoylase (PyrI), suggesting that either a paralogous gene serves this function or PyrB functions in the absence of a regulatory subunit.

*H. pylori* has homologues for all of the enzymes used for the interconversion of pyrimidine deoxyribonucleotides (Fig. 3B) with the exception of thymidylate synthase (ThyA), which is required for the interconversion of dUMP to dTMP. In addition, homologues for all of the enzymes associated with the pyrimidine salvage pathway are absent except for DeoB. This absence is consistent with poor utilization of uracil and uridine and with the failure to detect incorporation of added thymine, cytosine, or deoxycytidine into DNA by *H. pylori* (108). Although thymidine kinase activity has been found in crude extracts of *H. pylori* (108), no gene encoding a homologue of thymidine kinase (*tdk*) was identified in *H. pylori*.

#### Inorganic Elements and Heavy Metals

Phosphorus is an essential element in bacteria. *H. pylori* possesses homologues to both polyphosphate kinase and inorganic pyrophosphatase. These enzymes confer the ability to synthesize and hydrolyze polyphosphate, in agreement with experimental evidence (17). *H. pylori* also possesses a phosphate transporter (JHP1384/HP1491).

Sulfur assimilation is restricted in *H. pylori* compared to *E. coli*. Homologues to the genes necessary for the assimilation of sulfide and cysteine (*cysE* and *cysK*) are present in both sequenced strains, whereas those for the assimilation of sulfate (*cysA*, *cysC*, *cysD*, *cysH*, and *cysN*), an energy-consuming process, are not. The absence of an identifiable sulfate permease supports the apparent inability of *H. pylori* to use sulfate. Whereas *H. pylori* can utilize only sulfide as a source of inorganic sulfur, the closely related bacterium *Campylobacter jejuni* has homologues to the genes necessary to assimilate sulfate, sulfite, and sulfide (based on analysis of the recently completed genome by the Sanger Centre). This difference in sulfur assimilation between *H. pylori* and *Campylobacter* spp. is also seen in sulfur dissimilation. Unlike many *Campylobacter* spp., *H. pylori* does not have the homologues necessary for the respiration of many sulfur compounds. The absence of these sulfur assimilatory and dissimilatory genes in *H. pylori* probably reflects the evolved physiology resulting from its unique gastric niche.

In *E. coli* and *Salmonella typhimurium*, nitrogen is derived mainly from the primary products of ammonia assimilation, i.e., glutamate and glutamine. Glutamine synthetase (GlnA)

TABLE 4. Predicted biosynthetic abilities and auxotrophies of *H. pylori*

Category and Compound	No. of genes present <sup>a</sup>	Predicted synthetic ability <sup>b</sup>
<b>Amino acids</b>		
Aspartic acid	2 (2)	Y
Cysteine	2 (3) <sup>c</sup>	Y
Glutamic acid	1 (1)	Y
Glutamine	1 (1)	Y
Glycine	1 (1)	Y
Lysine	6 (7) <sup>d</sup>	Y
Threonine	5 (5)	Y
Tryptophan	5 (5)	Y
Alanine	1 (2)	N
Arginine	0 (9)	N
Histidine	0 (8)	N
Isoleucine	2 (9)	N
Leucine	1 (6)	N
Methionine	1 (7)	N
Phenylalanine	1 (3)	N
Proline	1 (3)	N
Valine	2 (9)	N
Asparagine	0 (2) <sup>e</sup>	?
Serine	2 (3)	?
Tyrosine	1 (3)	?
<b>Cofactors and vitamins</b>		
Biotin	4 (4)	Y
CoA	0 (5) <sup>d</sup>	Y
Folate	7 (10) <sup>d</sup>	Y
Molybdopterin	8 (9) <sup>e</sup>	Y
Panthenate	4 (4)	Y
Protoheme	10 (10)	Y
Pyridine nucleotides	3 (5) <sup>d</sup>	Y
Pyridoxal phosphate	3 (3) <sup>d</sup>	Y
Riboflavin	6 (7) <sup>e</sup>	Y
Thiamine	3 (9) <sup>e,d</sup>	Y
Thioredoxin	2 (2)	Y
B <sub>12</sub>		
Glutathione	0 (2)	N
Siroheme	0 (1)	N
Ubiquinone	3 (8)	?
Menaquinone	0 (6)	? <sup>f</sup>
<b>Polyamines</b>		
Agmatine	1 (1)	Y
Putrescine	1 (3)	N
Spermidine	3 (5)	? <sup>g</sup>
Pyrimidines	11 (10) <sup>h</sup>	Y
Purines	3 (10)	N <sup>i</sup>

<sup>a</sup> Number of genes assigned in *E. coli* shown in parentheses. Data from reference 119.

<sup>b</sup> Y, yes; N, no; ?, not clear.

<sup>c</sup> *H. pylori* has a gene for each step in the biosynthetic pathway. The number of genes in *E. coli* is larger due to redundancy.

<sup>d</sup> Not all genes in this pathway have been identified in *E. coli*.

<sup>e</sup> Synthesis may occur via tRNA (see the text).

<sup>f</sup> The presence of menaquinones in *H. pylori* has been demonstrated (see the text).

<sup>g</sup> Spermidine synthesis may be mediated by NapC.

<sup>h</sup> Two copies of *pyrC* were found in *H. pylori*.

<sup>i</sup> Salvage pathway is present (see the text).

catalyzes the formation of glutamine, while either glutamate dehydrogenase (GdhA) or glutamate synthase (GltBD) catalyzes the formation of glutamate (131). Homologues to *glnA* and *gdhA* were identified, and the gene product of *glnA* has been characterized in *H. pylori* (45). Glutamate dehydrogenase







succinate, and lactate (99, 106), an experimental observation consistent with the genomic analyses.

**Fermentation.** *H. pylori* ferments pyruvate to acetate. Genetic analysis indicates that strain J99 can carry out this fermentation but that strain 26695 cannot do so due to a frame-shift in its phosphotransacetylase gene. This mutation also implies that strain 26695 cannot convert acetate to acetyl-CoA by running the fermentative pathway in the reverse direction. However, in *H. pylori* 26695 the single, identifiable strain-specific gene involved in energy metabolism is an acetyl-CoA synthetase (HP1045) (149), which allows the direct conversion of acetate to acetyl-CoA.

The presence of alcohol dehydrogenase activity in *H. pylori* suggests that this bacterium can ferment pyruvate to ethanol (135, 136). Indeed, a homologue of this enzyme was identified in both strains. In addition, the single, identifiable strain-specific gene involved in energy metabolism in strain J99 (JHP1429) is a second alcohol dehydrogenase homologue. The role of alcohol dehydrogenase in *H. pylori* is unclear, because this organism apparently cannot ferment pyruvate to ethanol due to the absence of an identifiable acetaldehyde dehydrogenase homologue. The absence of such a homologue is consistent with biochemical studies (135, 136). The ability of *H. pylori* to produce acetaldehyde from ethanol (133) suggests that this bacterium is very sensitive to alcohols since it cannot detoxify the resulting acetaldehyde.

**Tricarboxylic acid cycle.** The tricarboxylic acid (TCA) cycle of *H. pylori* in both sequenced strains is similar to the branched anaerobic TCA pathway used by *E. coli* (Fig. 2) with the following exceptions. Succinyl-CoA is generated from 2-oxoglutarate rather than from succinate (64). Furthermore, in *H. pylori*, fumarate is generated in the TCA pathway from aspartate rather than from malate. In contrast, genomic analysis suggests that in the closely related bacterium *C. jejuni*, succinyl-CoA is synthesized from succinate and that fumarate is synthesized from malate (based on analysis of the recently completed genome by the Sanger Centre). *H. pylori* possesses a homologue of fumarase, explaining the conversion of malate to fumarate observed in crude extracts (101). Unlike *C. jejuni*, no other malate-utilizing enzymes, such as malate dehydrogenase, malate synthase, or malate oxidoreductase, were identified in *H. pylori*.

**Fatty acid degradation.** Both sequenced strains of *H. pylori* contain the genes necessary for C<sub>2</sub> or short-chain fatty acid catabolism (25) and for a short-chain fatty acid transporter. No identifiable homologues were found to the genes involved in long-chain fatty acid  $\beta$ -oxidation. Together, these observations indicate that *H. pylori* may utilize acetoacetate and not acetobutyrate as a source for short chain fatty acid catabolites.

### Electron Transport Chain

**Electron donors.** The initial transfer of electrons during the oxidation of D-lactate and NADH may be performed by homologues to D-lactate dehydrogenase (*dld*), NADH dehydrogenase I (*nuoA-nouN*), and hydrogenase (*hyaA* to *hyaDD*). The presence of these genes in both sequenced strains is consistent with measured activities in *H. pylori* cell membranes (22, 93, 101, 105). Although homologues to NuoE and NuoF were not identified in *H. pylori*, the *nuo* cluster contained two ORFs in an identical location and of similar size to the two *E. coli* genes. Thus, it is likely that these two *H. pylori* ORFs encode proteins with orthologous functions to NuoE and NuoF. In addition, homologues for the following electron-transferring systems were found: pyruvate ferredoxin oxidoreductase (*porA*, *porB*,

*porG*, and *porD*), glycerol-3-phosphate dehydrogenase (*glpC*), and proline dehydrogenase (*putA*).

No genes encoding a succinate dehydrogenase homologue were identified, although such an activity has been observed in extracts from several *H. pylori* strains, including J99 (34, 59, 120). This apparent discrepancy can be explained by the observation that fumarate reductase can convert succinate to fumarate in vitro. The importance of fumarate reductase in respiration depends on environmental conditions. Under microaerophilic conditions, fumarate reductase (46) is not essential in *H. pylori*, which explains why high concentrations of fumarate reductase-specific antimicrobials are required to inhibit the growth of and kill *H. pylori* in vitro (105). Under these conditions, oxygenic respiration may be used by the bacterium and hence reduces the importance of fumarate reductase in metabolism. In the absence of oxygen, this enzyme may be essential. In the presence of oxygen and fumarate, *H. pylori*, like members of the related genus *Wolinella*, may prefer fumarate as a terminal electron acceptor over performing oxygenic respiration (61). The role fumarate reductase plays in respiration cannot be assessed until the microenvironment of *H. pylori* is better defined.

**Quinones and cytochromes.** Analysis of the genomic sequence provides no clear indication to the composition of the *H. pylori* quinone pool. No homologues to genes involved in the biosynthesis of menaquinones, elements of anaerobic respiration, were identified despite the reported presence of menaquinone-6 and menaquinone-1 in the cell membranes of *H. pylori* (52, 94). The absence of identifiable homologues implies that *H. pylori* obtains menaquinones either by synthesis with genes that have yet to be identified or by uptake from its environment. *H. pylori* does contain homologues for three ubiquinone-biosynthetic enzymes (UbiA, UbiD, and UbiE), but no significant homologues for UbiB, UbiC, UbiF, UbiG, or UbiH were identified. UbiB, UbiF, and UbiH are oxygen-utilizing enzymes, and their apparent absence in *H. pylori* might be the result of the microaerophilic metabolism of the bacterium.

The biosynthesis of cytochromes by *H. pylori* has been reviewed in detail recently (51, 125). *H. pylori* uses a type II system for such biosynthesis, which is similar to that in many gram-positive bacteria and some members of the  $\beta$ -proteobacteria. Both strains possess homologues to all of the components needed for this biosynthetic system.

**Terminal electron acceptors.** There appear to be three putative terminal electron acceptor systems in both sequenced strains of *H. pylori*, i.e., fumarate reductase, *N*-oxide reductase, and cytochrome *c* oxidase, suggesting that *H. pylori* may be able to use fumarate, *N*-oxides (i.e., dimethyl sulfoxide and trimethylamine-*N*-oxide), or oxygen as electron sinks. Whereas the presence of a cytochrome *c* oxidase in *H. pylori* would allow aerobic respiration, the presence of both fumarate reductase and an *N*-oxide reductase suggests that the bacterium may respire anaerobically as well. The terminal oxidase complex is similar to *cbb*<sub>3</sub>-type oxidase complexes and is encoded by a gene cluster composed of homologues to genes encoding the *Rhizobium* FixN, FixO, FixP, and FixQ subunits. The arrangement of these genes is identical to that found in *Rhizobium* spp. (130), and the presence of such a terminal oxidase in *H. pylori* is consistent with previous findings (116).

### ATP-Proton Motive Force Conversion

The bacterial ATP synthase, a multisubunit enzyme, is composed of the F<sub>0</sub> complex, which consists of three subunits that form a proton channel, and the F<sub>1</sub> complex, which consists of

five subunits that constitute the catalytic site for ATP synthesis. In *E. coli*, all eight subunits are encoded within the *atp* operon (76). All five subunits of the F<sub>1</sub> complex and the F<sub>0</sub> b subunit are contiguous on the *H. pylori* chromosome. The remaining two subunits of the F<sub>0</sub> complex are encoded by genes present in other chromosomal regions. *H. pylori* has an additional subunit (JHP1065/HP1137), which is homologous to the ATPase b', a diverged and duplicated form of the b subunit found among plants and photosynthetic bacteria. The gene encoding this homologue is located at one end of the ATP synthase gene cluster. Functionally, the *H. pylori* ATPase is similar to other bacterial ATPases in that it uses the proton motive force generated by the electron transport chain to synthesize ATP (97).

### Detoxification

Organisms that come in contact with oxygen, like the microaerophilic *H. pylori*, must be able to protect themselves from the toxic products of oxygen metabolism, such as superoxide and hydrogen peroxide. Both sequenced strains of *H. pylori* possess a superoxide dismutase and a catalase, consistent with previous biochemical findings (57, 122, 140). Thus, *H. pylori* is a microaerophile not because of an inability to neutralize the toxic products of oxygen metabolism but, more probably, as a consequence of other metabolic limitations. *H. pylori* has two genes encoding peroxidases (JHP991/HP0390 and JHP1471/HP1563), one of which is located adjacent to the superoxide dismutase gene. The ability to isolate catalase-negative mutants of *H. pylori* (122, 156) suggests that at least one of the peroxidases can function as a catalase.

In summary, it would appear that *H. pylori* can use amino acids or simple carbohydrates as a major source of energy. The bacterium is restricted with respect to pyruvate metabolism. Further, *H. pylori* possesses an electron transport chain that can use oxygen as a terminal electron acceptor, but homologues to fumarate reductase and *N*-oxide reductase suggest that the bacterium is capable of at least limited anaerobic metabolism.

## MACROMOLECULE BIOSYNTHESIS AND MODIFICATION

### DNA Replication, Recombination, and Restriction-Modification

*H. pylori* contains genes encoding homologues to the DnaE, DnaN, DnaX, DnaQ, and HolB subunits of the DNA polymerase III holoenzyme, which is responsible for DNA replication. While the *H. pylori* holoenzyme contains fewer than the 10 subunits that comprise DNA polymerase III in *E. coli*, the total number of subunits is consistent with that found in other bacterial genomes (24, 31, 43, 81, 84, 139). Indeed, the only common subunits among these different species are DnaE, DnaN, and DnaX. *H. pylori* contains homologues to all genes encoding enzymes involved in initiation and DNA chain elongation, except *dnaC*, which is also absent in several other bacterial genomes (24, 31, 43, 81, 84, 139).

*H. pylori* possesses homologues to several nucleases, including UvrABC endonuclease, UvrD, ExoA, and RecJ. Even though *H. pylori* contains homologues to *xseA* and *mutS*, it lacks recognizable homologues encoding the other subunits of exonuclease VII (XseB) and the MutHLS endonuclease repair complex, respectively. Recombinational repair in *H. pylori* appears to occur in a RecBC-independent manner (92), since the

RecBCD exonuclease V is absent and a RecR homologue was identified. Other homologues identified in the recombination system include RecA and RecN, while no homologue to *recE* encoded exonuclease VIII was found. In addition, homologues of *ruvABC* and *recG*, whose products are involved in the branch migration and resolution of Holliday structures, were identified in *H. pylori*. Both *H. pylori* genomes also possess several homologues to the DNA topoisomerase I gene (*topA*), some of which are strain specific (6). There are genes encoding several other ATP-dependent helicases in *H. pylori*, including a homologue for *pcrA* and *rep*. Both sequenced *H. pylori* strains have numerous genes with similarity to DNA restriction and modification enzymes, many of which are strain specific. This finding suggests that *H. pylori* strains have their own unique complement of these genes (2, 6). While many of the products of these genes can be classified as type I, II, or III restriction or modification enzymes, their exact specificity remains to be determined.

Both *H. pylori* genomes appear to have three type I systems. In all three systems, the modification (HsdM) and restriction (HsdR) subunits are highly homologous between the strains but the specificity subunits (HsdS) have limited identity. Domains of HsdS proteins can be shuffled to produce new specificity, and it is this modular nature which allows the type I systems to evolve rapidly to a new DNA specificity.

Some of the type II systems in *H. pylori* 26695 and J99 appear to be functionally equivalent and may possess the same DNA specificity, while others have identity within the methyltransferase enzyme but differences in the restriction enzyme. There are nine type II modification methylases common to both genomes (6). In addition, *H. pylori* J99 possesses two unique type II restriction-modification systems. In *H. pylori* 26695 and several other strains, the *M.HpyI* gene, encoding a type II modification enzyme (159), flanks *iceA*, which encodes a putative type II restriction endonuclease (149). The absence of *iceA* in *H. pylori* J99, a recent clinical isolate from a patient diagnosed with a duodenal ulcer, suggests that *iceA* is not required for gastrointestinal disease and may not represent an informative epidemiological marker of pathogenicity and virulence, as previously hypothesized (15, 152).

Both *H. pylori* 26695 and J99 contain two type III restriction-modification systems, one of which is strain specific. Whereas in the other system (JHP1410/1411, HP1521/1522) the restriction gene product is 93% identical in the two strains, the modification gene product is conserved only at the N and C termini. This difference may result in unique specificity for each modification enzyme. The modification gene appears to be regulated by a slipped-strand repair mechanism and is frameshifted in both J99 and 26695 (6).

### Transcription and Translation

Analysis of the *H. pylori* genomes identified only three sigma factors (RpoD, RpoN, and FliA). The presence of these three sigma factors had been suggested by putative promoters found upstream of *H. pylori* genes (83, 86, 89, 122, 141, 147). No homologue to the stationary-phase sigma factor (RpoS) or the heat shock-specific sigma factor (RpoH) was identified, implying that *H. pylori* responds to stress in different fashion from that described in other bacteria. Both sequenced *H. pylori* strains contain a fusion of *rpoB* and *rpoC*, which encode the  $\beta$  and  $\beta'$  subunits of RNA polymerase (149, 160). *H. pylori* contains homologues to three termination factors (*nusA*, *nusB*, and *rho*) and lacks identifiable homologues to the tRNA maturation genes *md*, *rph*, and *mpB*. The lack of identifiable tran-






FIG. 4. Peptidoglycan synthesis and recycling. GlcNAc, *N*-acetylglucosamine; MurNAc, *N*-acetylmuramic acid; Ala, alanine; Glu, glutamate; Dap, diaminopimelic acid. Adapted from reference 62.

scriptional termination stem-loop structures suggests that in *H. pylori* termination is largely Rho dependent (6).

All the aminoacyl-tRNA synthetases are present in *H. pylori*, except glutamyl- and asparaginyl-tRNA synthetases. Two copies of the gene encoding glutamyl-tRNA synthetase (*gltX*) are present in both *H. pylori* strains. It had been suggested that one of these copies may function as a glutamyl-tRNA synthetase (149). However, the presence of homologues to the *gatA*, *gatB*, and *gatC* genes, which have been demonstrated to replace glutamyl-tRNA synthetase activity in *Bacillus subtilis* (29, 146), makes it more likely that glutamyl-tRNA synthetase activity in *H. pylori* is encoded by the *gatABC* homologues rather than *gltX* (6). Thus, the role of the second glutamyl-tRNA synthetase in *H. pylori* is unclear. A similar transamidation reaction, encoded by these three homologues or other genes, may function in *H. pylori* to synthesize asparaginyl-tRNA from aspartyl-tRNA, explaining the ability of *H. pylori* to grow without added asparagine despite its apparent inability to synthesize this amino acid.

### Fatty Acid and Phospholipid

Identifiable homologues were found to many of the genes required for initiation and elongation of fatty acid biosynthesis. Both *H. pylori* strains contain the characteristically small acyl carrier protein (ACP) (78 amino acids) (JHP744/HP0808). *H. pylori* 26695 but not J99 possesses a second, significantly larger (153-amino-acid) homologue with an extended N-terminal domain (HP962). Genomic analysis indicates that *H. pylori* has a homologue to cyclopropane fatty acid synthase (JHP969/HP0416), consistent with the experimental evidence that *H. pylori* has a preponderance of C<sub>19:0</sub> cyclopropane chains (73). No homologue to  $\beta$ -hydroxydecanoyl-ACP dehydrase, which catalyzes the formation of *cis*-3-decenoyl-ACP, an important

intermediate in the biosynthesis of unsaturated fatty acids in *E. coli*, was found. The absence of this homologue is surprising, since unsaturated fatty acids are present in *H. pylori* (73).

The phospholipid composition of *H. pylori* consists mainly of phosphatidylethanolamine, cardiolipin, and phosphatidylglycerol, with smaller quantities of phosphatidylserine and phosphatidylcholine (73). The genome appears to encode all of the proteins necessary for the synthesis of these phospholipids except for cardiolipin synthase (*cls*), which catalyzes the final step of cardiolipin synthesis in *E. coli*. The *H. pylori* genome encodes at least two of the three enzymes necessary for phosphatidic acid synthesis (JHP895/HP0961 and JHP1267/HP1348). A homologue to the glycerol-3-phosphate acyltransferase was not identified in the *H. pylori* genome.

A characteristic feature of the lipid profile of *H. pylori* is the presence of cholesterol glucosides, which account for about 25% of the total lipid of the bacterium (56, 58). *H. pylori* does not appear to encode known enzymes for the synthesis of cholesterol and presumably scavenges this molecule from the environment. Little is known about glucoside-modified cholesterol synthesis, but the enzymes responsible are expected to be found among the species-specific genes with unknown function.

### Peptidoglycan

The cytoplasmic synthesis of UDP-activated precursors of bacterial peptidoglycan assembly is well understood. The *H. pylori* genome encodes homologues to all of the enzymes in this pathway, beginning with the synthesis of UDP-*N*-acetylmuramic acid and ending with UDP-disaccharide pentapeptide linked to an undecaprenol lipid carrier (Fig. 4).

After transport through the cytoplasmic membrane, peptidoglycan precursors are incorporated into the existing pepti-

doglycan layer by the penicillin-binding proteins (PBPs) (50). The genome of *H. pylori* encodes three proteins which show homology to PBPs, in agreement with a published gel electrophoresis study (68). The precise metabolic function of each PBP is uncertain. However, one of the PBPs (JHP544/HP597) has a transglycosylase motif. Since no additional genes with a similar motif were detected, this PBP may be the only enzyme in *H. pylori* that is involved in lengthening the glycan chain.

From the analysis of the genomes, it is uncertain if peptidoglycan fragments can be recycled by *H. pylori*. Although this bacterium has homologues to genes encoding a lytic amidase (JHP709/HP0772) and an *N*-acetylmuranoyl-*L*-alanine transglycosylase (JHP590/HP0645), it does not appear to have the other genes required for this recycling system (*ampG*, *ampD*, *mpl*) (127).

### Outer Membrane

Approximately 4% of the coding capacity of both strains is devoted to outer membrane proteins. This amount is significantly larger than that of any other bacterial genome sequenced to date. The majority of these proteins belong to three paralogous families, the largest having 20 and 21 members in J99 and 26695, respectively (6). Several members of the largest paralogous family are porins (33, 41) or adhesins specific for the Lewis B carbohydrate moiety found on host cells (72). The sequence identity of orthologous members of this large family is high (greater than 95%), suggesting that strain-specific sequence differences play only a limited role in antigenic variation. Five orthologous pairs of this outer membrane protein family, including BabB (Lewis B adhesin), contain dinucleotide (CT) repeats in their signal sequences. Slipped-strand repair has been proposed to regulate the expression of these proteins (6, 149). Significantly, the predicted expression status (in frame or out of frame) of these five pairs is identical despite the presence of a different number of dinucleotide repeats in the two strains in each case (6).

### Lipopolysaccharide

*H. pylori* has homologues to all enzymes required for 2-keto-3-deoxyoctulosonic acid (KDO)-lipid A biosynthesis. Compared to lipid A in *E. coli*, this moiety in *H. pylori* is underacylated, has longer fatty acid chains ( $C_{16}$  and  $C_{18}$ ), and lacks a phosphate group at the 4'-hydroxyl position on the nonreducing glucosamine of the disaccharide (115). Taken together, these findings suggest that in *H. pylori* lipid A is assembled as an acylated and diphosphorylated disaccharide, which is then modified by an unidentified phosphatase and esterase enzyme(s).

The chemical structures of the lipopolysaccharide cores from two *H. pylori* strains were shown to be identical heptasaccharides (9, 10). Synthesis of such a structure requires several glycosyltransferases. There are seven ORFs encoding these putative glycosyltransferases (JHP147/HP0159, JHP194/HP0208, JHP563/HP0619, JHP620/HP0679, JHP741/HP0805, JHP765/HP0826, and JHP1031/HP1105), which are common to both strains, three strain-specific ORFs in J99 (JHP562, JHP820, and JHP1032), and one strain-specific ORF in 26695 (HP1578). However, none of these ORFs can be assigned a substrate specificity. Whether the presence of these strain-specific glycosyltransferases in J99 and 26695 results in a different lipopolysaccharide core structure remains to be determined.

The O chain from lipopolysaccharides of *H. pylori* is composed of Lewis acids [ $Le^x$  and  $Le^y$ ; mono- and difucosylated repeating disaccharides of  $\beta$ -(1,4)-linked galactose and *N*-acetylglucosamine, respectively] (9, 10). These carbohydrate

moieties, which are identical to those found on host tissues, have been implicated in colonization and persistence and may also play a role in autoimmunity (7, 8). The biosynthetic pathway of these O chains has not been determined. One enzyme,  $\alpha$ -(1,3)-fucosyltransferase (JHP596/HP0651), is to be involved in this pathway (21, 96). Each genome has two  $\alpha$ -(1,3)-fucosyltransferases which differ in the number of a 7-amino-acid sequence repeat (YDDLRVN). Regulation of these genes appears to occur through slipped-strand repair at two distinct polynucleotide repeats (6).

### Flagella

Flagellar biosynthesis in gram-negative bacteria has been extensively studied. The assembly of a functional flagellum requires numerous proteins and is a highly regulated process. Homologues to all of the required genes involved in flagellar biosynthesis were identified in both sequenced strains. In other gram-negative bacteria, including *C. jejuni*, inactivation of the biosynthetic pathway disrupts the expression of the system as a whole. However, in *H. pylori*, inactivation of the hook protein does not result in suppression of flagellin expression (124), indicating that flagellar biosynthesis is not as highly regulated as in other bacteria. This difference can be explained by the absence of FlgM in *H. pylori*, a protein that in other bacteria controls feedback regulation of the flagellar biosynthetic cascade.

The flagellar filament of *H. pylori* is composed of two flagellin subunits, FlaA and FlaB, and genes encoding both are present. Despite reports suggesting that the genes for these flagellin subunits exhibit strain variation (66, 123), in strains J99 and 26695 the FlaA protein sequences are identical and the FlaB protein sequences differ by only a single amino acid. In addition, the protein sequences reported for FlaA and FlaB from another strain are nearly identical to those from strains J99 and 26695 (89).

The flagellar filament of *H. pylori* is encased within a sheath that is continuous with the outer membrane (90, 91, 128). The sheath, whose composition has yet to be defined, probably protects the polymeric filament from dissociation in the low pH of the stomach. Likewise, other polymeric structures, such as fimbriae, would also be subject to dissociation at low pH, thus explaining why adhesion to the host epithelium by *H. pylori* appears to be mediated by integral outer membrane proteins (71, 72).

In summary, *H. pylori* has the necessary homologues for DNA, RNA, and protein synthesis.

## CELLULAR PROCESSES

### Protein Secretion

*H. pylori* has homologues to two leader peptidases, enzymes required for protein secretion. The type I leader peptidase, LepB, is responsible for cleavage of the signal sequence from most periplasmic and outer membrane proteins. The *lspA* (previously called *ureD*) gene product is the leader peptidase responsible for processing prelipoproteins.

The secretion of proteins through the cytoplasmic membrane utilizes specific machinery which consists of several *sec* gene products. *H. pylori* contains all the known Sec proteins except SecB and SecG. Mutation of *secB*, which encodes a chaperone, affects the secretion of only a limited number of proteins in *E. coli*, and other preproteins may utilize alternative chaperones. Indeed, it has been demonstrated that *groESL* mutants in *E. coli* have significant effects on the secretion of

some proteins (85). Thus, it is likely that in *H. pylori*, other cytoplasmic chaperones are used to usher proteins, in conjunction with SecA, to the cytoplasmic membrane in preparation for secretion. SecE is proposed to be integral to the translocation machinery (121). Initial analysis of both genomes did not identify a SecE orthology. However, a region between JHP1126/HP1203 and JHP1127/HP1204 contains a 59-amino-acid ORF that has similarity to the functionally important region of *E. coli* SecE.

### Cag Pathogenicity Island

*H. pylori* strains associated with clinically severe gastric disease (peptic ulcers) more commonly possess an approximately 40-kb pathogenicity island containing cytotoxin-associated genes (Cag pathogenicity island [cagPAI]) than do strains isolated from patients with uncomplicated gastritis (26). Type I strains have been defined as *H. pylori* isolates which have the entire cagPAI, express the cytotoxicity-associated immunodominant antigen (CagA) and an active vacuolating cytotoxin (VacA), and induce interleukin-8 (IL-8) secretion by gastric epithelial cells. In addition to cagA, it has been demonstrated by mutational analysis that several of the genes in the cagPAI are required for wild-type induction of IL-8 secretion by gastric epithelial cells. In contrast, Type II strains do not express CagA, have no vacuolating activity even though a truncated VacA may be produced, and do not induce IL-8 secretion at levels comparable to that induced by type I strains.

The initial cagPAI sequence was produced in two parts by two different laboratories, each of which sequenced clones from the same ordered cosmid library that had been constructed from *H. pylori* NCTC 11638 (2, 19). These groups showed that the cagPAI was separated into the cagI and cagII segments by an intervening stretch of DNA that was itself bordered on each side by a newly identified insertion sequence (IS605) element. In contrast, the cagPAI in *H. pylori* J99 and 26695, both type I strains, consists of the cagI and cagII segments fused as a single unit without the stretch of intervening DNA flanked on each side by IS605. In all three strains, the cagI and cagII segments are essentially the same and the cagPAI is found in the same relative location.

Many of the genes in the cagPAI are required for specific host cell responses to infection by *H. pylori*, including induction of (i) IL-8 secretion by gastric epithelial cells (2, 19, 137), (ii) tyrosine phosphorylation in host proteins (137), and (iii) cytoskeletal rearrangements during actin pedestal formation at the host cell surface (137). The manner in which cagPAI genes cause these host cell responses is unknown. However, five ORFs show significant sequence similarity to genes encoding the VirB protein family, which is responsible for DNA transfer in *Agrobacterium tumefaciens* (type IV secretion system) (23). These genes also have sequence similarity to genes involved in conjugative transfer of plasmids in *E. coli* and protein export in *Bordetella pertussis* (155). It is unknown how these components of conjugative and protein transport systems function in *H. pylori*. It has been suggested that the cagPAI encodes a type of contact-mediated secretion apparatus analogous to the type III secretion systems identified and characterized in several enteric pathogens such as *Yersinia*, *Salmonella*, *Shigella*, and pathogenic *E. coli* (88). It is therefore likely that the virB homologues of the cagPAI region encode components of the secretion apparatus which may deliver effector molecules (DNA or protein) directly to the host cell to elicit the responses mentioned above.

### Insertion Elements

The IS element IS605 contains genes encoding two previously identified transposases flanked by a short nucleotide sequence with dyadic symmetry and a common central core sequence (19). IS605 transposes as a single unit in *E. coli*, suggesting that it could also be functional in *H. pylori* (79). One of the transposase genes (*tmpA*) is related to IS200 found in gram-negative bacteria, and the other (*tmpB*) is related to IS1341 found in a gram-positive thermophilic bacterium. It is unusual to find an insertion element with transposases from apparently two different origins. The IS605 element was first described within the cagPAI of NCTC 11638. Strain 26695 has five full copies of IS605 and one copy of a related insertion element, IS606, none of which is located within the cagPAI. Strain J99 has no complete copies of IS605 but has one copy of IS606 on its chromosome. The short flanking sequence of IS605, without the transposases (is605), is present on both ends of the cagPAI in strain NCTC 11638. These sequences flanking cagPAI are thought to be remnants of a recent transposition or of another type of recombinational event. The dyadic repeats of IS605 and IS606 are also found within the J99 and 26695 genomes, at both common and distinct locations. IS605 dyadic repeats are coincident, in one or the other genomes, with several of the major organizational differences between the two sequenced strains. *H. pylori* plasmid pHPM186 (GenBank accession no. AF077006) contains an IS605 element which is adjacent to three genes that flank an IS605 element in the plasticity zone of strain 26695 (6). This finding suggests that plasmid integration plays a role in generating the limited genomic diversity in *H. pylori*. Thus, insertion elements, such as IS605, may have been involved in the acquisition of cagPAI and the plasticity zone by *H. pylori* via horizontal transfer, as has also been postulated for other pathogenicity islands found in a wide range of pathogenic bacteria.

### Transformation

Many *H. pylori* strains are naturally competent for DNA transformation, and the efficiency of this process varies from strain to strain. Besides the conjugative DNA transfer/protein export homologues (VirB proteins) encoded by the cagPAI, both strains contain additional members of this family, some of which are strain specific. Recently, some members of the VirB family have been shown to play a role in DNA transformation in *H. pylori* (60). The existence of strain-specific VirB family homologues may explain the variation in DNA transformation efficiency seen between strains. One additional transformation-associated gene (*comEC*), which is required for uptake of DNA into *B. subtilis*, was also identified.

### Chemotaxis

Chemotaxis, the sensory adaptation mechanism by which motile bacteria recognize and react to environmental conditions, has been found in *H. pylori* (110). Three homologues of the chemotaxis pathway in *E. coli* (CheW, CheA, and CheY), as well as four methyl-accepting chemotaxis proteins (MCPs), which mediate specificity for ligands, were identified in both sequenced strains of *H. pylori*. Proteins with similarity to MCPs are not necessarily involved in flagellar chemotaxis (5, 30). Neither strain contains identifiable homologues to CheR or CheB, enzymes which, respectively, add methyl groups to or remove methyl groups from the MCPs, precisely modulating the chemotactic response. By contrast, *C. jejuni* does possess homologues to both CheR and CheB. Therefore, the chemotaxis observed in *H. pylori* may occur by a CheB- and CheR-



independent mechanism, similar to that seen in CheB CheR mutants of *E. coli* (145). The apparent inability of *H. pylori* to precisely modulate chemotaxis may reflect its limited but unique gastric niche.

Both *H. pylori* strains have three homologues to CheV, another chemotaxis protein, which has an N-terminal domain similar to CheW and a C-terminal response regulator domain similar to CheY (44). The CheW protein and the N-terminal domain of CheV are both capable of modulating the CheA-MCP interaction (134). The three CheV orthologous pairs have greater than 97% identity, whereas none of the paralogues have more than 40% identity at the amino acid level, suggesting that each orthologous pair has functional similarity and that each paralogue has a different specific function.

### Cell Division and Morphology

Several genes implicated in bacterial cell division were found in both sequenced strains of *H. pylori*. Among these are *ftsZ* and *ftsA*, which are adjacent to each other, as is generally observed (154). *H. pylori* does not possess an unambiguous homologue to FtsW, but it does possess two homologues to RodA, a protein with significant sequence similarity to FtsW (69). It is possible that one of the RodA homologues actually functions as FtsW. In *E. coli*, at least one of the peptidoglycan-synthesizing enzymes is specifically involved in cell division (142). *H. pylori* has three homologues to peptidoglycan-synthesizing enzymes, one of which could act specifically during cell division. Two homologues of the metalloprotease FtsH, phenotypically associated with cell division, were identified. Interestingly, mutagenesis studies have shown that one of the FtsH homologues (JHP356/HP1069) is essential for growth in vitro (48). This suggests that the second homologue (JHP271/HP0286) is unable to functionally replace the first homologue and probably has a different function. In addition, homologues of *ftsK* and *fic* were identified.

No homologues to ZipA, which is believed to initiate formation of the FtsZ ring (53), Sula, which is thought to inhibit the formation of the FtsZ ring (14), FtsQ, FtsN, or FtsL were identified. Also, no homologue to MinC, a cell division inhibitor, was found, whereas homologues to the activator (MinD) and cofactor (MinE) of MinC were identified.

Two homologues of cell morphology-determining proteins, RodA and MreB, were found. RodA is required for the catalytic activity of a PBP during elongation in *E. coli* (74), whereas the function of MreB is unknown.

### Virulence Factors

A number of *H. pylori* proteins have been implicated in pathogenesis (for reviews, see references 36 and 95). All of the reported genes encoding potential virulence factors in *H. pylori* were identified in both strains, with the exception of *iceA*, which is missing from strain J99 (see "DNA replication, recombination, and restriction-modification" above).

Bacterial motility has been suggested to be required for colonization of the gastric mucosa, since *H. pylori* mutants unable to synthesize either of the flagellar subunits (FlaA or FlaB) cannot colonize gnotobiotic piglets (38, 40). Whether motility is needed for the persistence of an infection is not known, but owing to the rapid gastric epithelium and mucous turnover, it would probably be required.

The cytotoxin VacA induces vacuole formation in cultured epithelial cells and may be an important component in the induction of gastric cell lesions by *H. pylori* (27, 49). Different alleles of the *vacA* gene have been reported (11, 28). Strains J99 and 26695 possess different *vacA* alleles (*s1b/ml* and *s1a/*

*ml*, respectively) and were isolated from patients with *H. pylori*-related disease of different severity. Epidemiological studies with humans have correlated the level of expression of the *vacA* gene with disease outcome (42), and particular *vacA* genotypes are associated with more severe disease symptoms (109, 144, 151). Interestingly, in gnotobiotic piglets, a mutant with a knockout mutation of *vacA* had no discernible effect on colonization, epithelial vacuolation, or gastritis, suggesting that VacA is not a virulence factor in this animal model (37).

The urease enzyme of *H. pylori* has been extensively studied (for reviews, see references 111 to 113) and has been shown to be a colonization factor (39, 78, 150). Urease is found in the cytoplasm as well as on the surface of *H. pylori*. The mechanism by which this enzyme is translocated is controversial (129, 153).

Other components which have been implicated in virulence, include the *cagPAI*, lipopolysaccharide, outer membrane proteins, and a number of enzymes, such as phospholipases, catalase, superoxide dismutase, and a mucinase homologue (36). The involvement of these components in pathogenesis remains to be elucidated. Homologues to the genes encoding all these products were present in both strains.

In summary, *H. pylori* possess homologues to the genes necessary for DNA replication, transcription, translation, and cell division, as well for as the synthesis of other important cellular macromolecules such as lipids and peptidoglycan. Each strain appears to possess a number of unique restriction enzymes that would result in selectivity to transformation with foreign DNA. Further features of this bacterium are the large number of related outer membrane proteins and the presence of a putative pathogenicity island.

### CONCLUSION

The genomic analysis suggests that both strains have essentially identical metabolic potential. The strain-specific genes that encode proteins with an assigned function are predicted to have little impact on the physiology of *H. pylori*. For the most part, the reported experimental data for biochemical activities present or absent in *H. pylori* are in agreement with the predicted metabolic capabilities. Our genomic analysis reveals nutritional requirements which would restrict the environments in which *H. pylori* can survive and, in part, explains its limited niche. *H. pylori* has broad catabolic capabilities. While it can perform oxygenic respiration, anaerobic respiration, and fermentation, *H. pylori* is limited in what it can use as a carbon source. Carbohydrate utilization is restricted to glucose, via the Entner-Doudoroff pathway, and to sugars with shorter carbon backbones than C<sub>6</sub>, for which transport systems are present. In addition, *H. pylori* possesses numerous transporter systems for the uptake of amino acids, which must be available in the gastric environment. Other than carbohydrates, *H. pylori* apparently uses amino acids as an important source of carbon which would yield less energy than using hexoses and could result in slow growth as occurs in vitro. Further, the deamination of amino acids, together with the action of urease, requires a mechanism to deal with increased levels of ammonia.

Colonization by *H. pylori* involves an interaction between the outer membrane of the bacterium and the gastric epithelium of the host. The outer membrane composition of *H. pylori* is unique in its protein content and lipopolysaccharide structure, which are consistent with the persistence of *H. pylori* in a restricted niche. Compared to other bacteria, *H. pylori* devotes a significantly higher percentage of its coding capacity to that of outer membrane proteins, further emphasizing the importance of these proteins. A number of genes involved in determining the composition of the outer membrane are differen-

tially regulated by slipped-strand repair (6). This differential regulation and strain-specific outer membrane-related genes may play a role in the severity of *H. pylori*-related disease and the ability of *H. pylori* to persist chronically in its host. The other strain-specific genes may also play a role in these aspects of pathogenesis. None of these strain-specific genes with an assigned function are predicted to have a significant impact on the pathophysiological capabilities of *H. pylori*, although those with an unassigned function may be important. However, there is evidence that host factors are involved in *H. pylori*-related diseases (35), and our genomic analyses suggest that these host factors may play a more significant role than was previously appreciated.

The analysis presented here, like previous analyses of other sequenced bacterial genomes, has found several biochemical pathways for which we were unable to identify all of the genes which should be present for that pathway to be functional. Some of these genes may not be present in the genome. In other cases, the biochemical activity may be present within the organism but the gene responsible for this activity may be unidentifiable by current in silico techniques. The apparent incompleteness of the same metabolic pathways in two unrelated strains may suggest that these pathways are functional in *H. pylori* and that the unidentified genes are different from previously described orthologues. These genes would be among the 40% of the genome to which no function has been assigned.

For the first time, the genomes of two strains from the same bacterial species have been compared (6). This publication provided an opportunity to begin defining the physiology of *H. pylori*, a globally important pathogen. Future genomic comparisons of multiple strains, carefully correlated with epidemiological data, will identify the minimal genomic complement of this species and the genes required for virulence. Such an approach will be applied to other pathogenic bacteria, and the genes identified from these studies will become candidates for therapeutic intervention.

#### REFERENCES

- Abouhamad, W. N., M. Manson, M. M. Gibson, and C. F. Higgins. 1991. Peptide transport and chemotaxis in *Escherichia coli* and *Salmonella typhimurium*: characterization of the dipeptide permease (Dpp) and the dipeptide-binding protein. *Mol. Microbiol.* **5**:1035–1047.
- Akopyants, N., S. Clifton, D. Kersulyte, J. Crabtree, B. Youree, C. Reece, N. Bukanov, E. Drazek, B. Roe, and D. Berg. 1998. Analyses of the *cag* pathogenicity island of *Helicobacter pylori*. *Mol. Microbiol.* **28**:37–53.
- Akopyanz, N., N. Bukanov, T. Westblom, and D. Berg. 1992. PCR-based RFLP analysis of DNA sequence diversity in the gastric pathogen *Helicobacter pylori*. *Nucleic Acids Res.* **20**:6221–6225.
- Akopyanz, N., N. Bukanov, T. Westblom, S. Kresovich, and D. Berg. 1992. DNA diversity among clinical isolates of *Helicobacter pylori* detected by PCR-based RAPD fingerprinting. *Nucleic Acids Res.* **20**:5137–5142.
- Alm, R., and P. Manning. 1990. Characterization of the *hlyB* gene and its role in the production of the El Tor haemolysin of *Vibrio cholerae* O1. *Mol. Microbiol.* **4**:413–425.
- Alm, R. A., L.-S. Lee, D. T. Moir, B. L. King, E. D. Brown, P. C. Doig, D. R. Smith, B. Noonan, B. C. Guild, B. L. deJonge, G. Carmel, P. J. Tummino, A. Caruso, M. Uria-Nickelsen, D. M. Mills, C. Ives, R. Gibson, D. Merberg, S. D. Mills, Q. Jiang, D. E. Taylor, G. F. Vovis, and T. J. Trust. 1999. Genomic sequence comparison of two unrelated isolates of the human gastric pathogen *Helicobacter pylori*. *Nature* **397**:176–180.
- Appelmelk, B., R. Negrini, A. Moran, and E. Kuipers. 1997. Molecular mimicry between *Helicobacter pylori* and the host. *Trends Microbiol.* **5**:70–73.
- Appelmelk, B., I. Simoons-Smit, R. Negrini, A. Moran, G. Aspinall, J. Forte, T. deVries, H. Quan, T. Verboom, J. Maaskant, P. Ghiara, E. Kuipers, E. Bloemena, T. Tadema, R. Townsend, K. Tyagarajan, J. C. Jr., M. Monteiro, A. Savio, and J. deGraaff. 1996. Potential role of molecular mimicry between *Helicobacter pylori* lipopolysaccharide and host Lewis blood group antigens in autoimmunity. *Infect. Immun.* **64**:2031–2040.
- Aspinall, G. O., and M. A. Monteiro. 1996. Lipopolysaccharides of *Helicobacter pylori* strains P466 and Mo19—structures of the O antigen and core oligosaccharide regions. *Biochemistry* **35**:2498–2504.
- Aspinall, G. O., M. A. Monteiro, H. Pang, E. J. Walsh, and A. P. Moran. 1996. Lipopolysaccharide of the *Helicobacter pylori* type strain NCTC 11637 (Atcc 43504)—structure of the O antigen chain and core oligosaccharide regions. *Biochemistry* **35**:2489–2497.
- Atherton, J., P. Cao, R. P. Jr., M. Tummuru, M. Blaser, and T. Cover. 1995. Mosaicism in vacuolating cytotoxin alleles of *Helicobacter pylori*. *J. Biol. Chem.* **270**:17771–17777.
- Berg, D., R. Gilman, J. Lelwala-Guruge, K. Srivastava, Y. Valdez, J. Watanabe, J. Miyagi, N. Akopyants, A. Ramirez-Ramos, T. Yoshiwara, S. Recavarren, and R. Leon-Barua. 1997. *Helicobacter pylori* populations in Peruvian patients. *Clin. Infect. Dis.* **25**:996–1002.
- Berg, D. E., P. S. Hoffman, B. J. Appelmelk, and J. G. Kusters. 1997. The *Helicobacter pylori* genome sequence: genetic factors for long life in the gastric mucosa. *Trends Microbiol.* **5**:468–474.
- Bi, E., and J. Lutkenhaus. 1993. Cell division inhibitors SulA and MinCD prevent formation of the FtsZ ring. *J. Bacteriol.* **175**:1118–1125.
- Blaser, M. 1997. Heterogeneity of *Helicobacter pylori*. *Eur. J. Gastroenterol. Hepatol.* **9**:S3–S6.
- Blaser, M. 1997. Not all *Helicobacter pylori* strains are created equal: should all be eliminated? *Lancet* **349**:1020–1022.
- Bode, G., F. Mauch, H. Ditschuneit, and P. Malfrather. 1993. Identification of structures containing polyphosphate in *Helicobacter pylori*. *J. Gen. Microbiol.* **139**:3029–3033.
- Brenchley, J. E., and B. Magasanik. 1974. Mutants of *Klebsiella aerogenes* lacking glutamate dehydrogenase. *J. Bacteriol.* **117**:544–550.
- Censini, S., C. Lange, Z. Xiang, J. Crabtree, P. Ghiara, M. Borodovsky, R. Rappuoli, and A. Covacci. 1996. *cag*, a pathogenicity island of *Helicobacter pylori*, encodes type I-specific and disease-associated virulence factors. *Proc. Natl. Acad. Sci. USA* **93**:14648–14653.
- Chalk, P. A., A. D. Roberts, and W. M. Blows. 1994. Metabolism of pyruvate and glucose by intact cells of *Helicobacter pylori* studied by <sup>13</sup>C NMR spectroscopy. *Microbiology* **140**:2085–2092.
- Chan, G. Z., M. M. Palcic, and D. E. Taylor. 1997. Cloning and heterologous expression of an  $\alpha(1,3)$ -fucosyltransferase from the gastric pathogen *Helicobacter pylori*. *J. Biol. Chem.* **272**:21357–21363.
- Chang, H. T., S. W. Marcelli, A. A. Davison, P. A. Chalk, R. K. Poole, and R. J. Miles. 1995. Kinetics of substrate oxidation by whole cells and cell membranes of *Helicobacter pylori*. *FEMS Microbiol. Lett.* **129**:33–38.
- Christie, P. J. 1997. *Agrobacterium tumefaciens* T-complex transport apparatus: a paradigm for a new family of multifunctional transporters in eubacteria. *J. Bacteriol.* **179**:3085–3094.
- Cole, S., R. Brosch, J. Parkhill, T. Garnier, C. Churcher, D. Harris, S. Gordon, K. Eglmeier, S. Gas, C. B. III, F. Tekaiia, K. Badcock, D. Basham, D. Brown, T. Chillingworth, R. Connor, R. Davies, K. Devlin, T. Feltham, S. Gentles, N. Hamlin, S. Holroyd, T. Hornsby, K. Jagels, A. Krogh, J. McLean, S. Moule, L. Murphy, K. Oliver, J. Osborne, M. Quail, M.-A. Rajandream, J. Rogers, S. Rutter, K. Seeger, J. Skelton, R. Squares, S. Squares, J. Sulston, K. Taylor, S. Whitehead, and B. Barrell. 1998. Deciphering the biology of *Mycobacterium tuberculosis* from the complete genome sequence. *Nature* **393**:537–544.
- Corthesy-Theulaz, I. E., G. E. Bergonzelli, H. Henry, D. Bachmann, D. F. Schorderet, A. L. Blum, and L. N. Ornston. 1997. Cloning and characterization of *Helicobacter pylori* succinyl CoA:acetoacetate CoA-transferase, a novel prokaryotic member of the CoA-transferase family. *J. Biol. Chem.* **272**:25659–25667.
- Covacci, A., S. Falkow, D. Berg, and R. Rappuoli. 1997. Did the inheritance of a pathogenicity island modify the virulence of *Helicobacter pylori*? *Trends Microbiol.* **5**:205–208.
- Cover, T., and M. Blaser. 1992. *Helicobacter pylori* and gastroduodenal disease. *Annu. Rev. Med.* **42**:135–145.
- Cover, T. L., M. K. Tummuru, P. Cao, S. A. Thompson, and M. J. Blaser. 1994. Divergence of genetic sequences for the vacuolating cytotoxin among *Helicobacter pylori* strains. *J. Biol. Chem.* **269**:10566–10573.
- Curnow, A., K. Hong, R. Yuan, S. Kim, O. Martins, W. Winkler, T. Henkin, and D. Soll. 1997. Glu-tRNA<sup>Gln</sup> amidotransferase: a novel heterotrimeric enzyme required for correct decoding of glutamine codons during translation. *Proc. Natl. Acad. Sci. USA* **94**:11819–11826.
- Darzens, A. 1994. Characterization of a *Pseudomonas aeruginosa* gene cluster involved in pilus biosynthesis and twitching motility: sequence similarity to the chemotaxis proteins of enterics and the gliding bacterium *Myxococcus xanthus*. *Mol. Microbiol.* **11**:137–153.
- Deckert, G., P. V. Warren, T. Gaasterland, W. G. Young, A. L. Lenox, D. E. Graham, R. Overbeek, M. A. Snead, M. Keller, M. Aujay, R. Huber, R. A. Feldman, J. M. Short, G. J. Olsen, and R. V. Swanson. 1998. The complete genome of the hyperthermophilic bacterium *Aquifex aeolicus*. *Nature* **392**:353–358.
- Dendinger, S. M., L. G. Patil, and J. E. Brenchley. 1980. *Salmonella typhimurium* mutants with altered glutamate dehydrogenase and glutamate synthase activities. *J. Bacteriol.* **141**:140–148.
- Doig, P., M. Exner, R. Hancock, and T. Trust. 1995. Isolation and characterization of a conserved porin protein from *Helicobacter pylori*. *J. Bacteriol.* **177**:5447–5452.



34. Doig, P., and T. Trust. 1994. Identification of surface-exposed outer membrane antigens of *Helicobacter pylori*. *Infect. Immun.* **62**:4526–4533.
35. Dubois, A., D. Berg, E. Incecik, N. Fiala, L. Heman-Ackah, G. Perez-Perez, and M. Blaser. 1996. Transient and persistent infection of nonhuman primates with *Helicobacter pylori*: implications for human disease. *Infect. Immun.* **64**:2885–2891.
36. Dunn, B., H. Cohen, and M. Blaser. 1997. *Helicobacter pylori*. *Clin. Microbiol. Rev.* **10**:720–741.
37. Eaton, K., T. Cover, M. Tummuru, M. Blaser, and S. Krakowka. 1997. Role of vacuolating cytotoxin in gastritis due to *Helicobacter pylori* in gnotobiotic piglets. *Infect. Immun.* **65**:3462–3464.
38. Eaton, K., D. Morgan, and S. Krakowka. 1992. Motility as a factor in the colonisation of gnotobiotic piglets by *Helicobacter pylori*. *J. Med. Microbiol.* **37**:123–127.
39. Eaton, K. A., C. L. Brooks, D. R. Morgan, and S. Krakowka. 1991. Essential role of urease in pathogenesis of gastritis induced by *Helicobacter pylori* in gnotobiotic piglets. *Infect. Immun.* **59**:2470–2475.
40. Eaton, K. A., S. Suerbaum, C. Josenhans, and S. Krakowka. 1996. Colonization of gnotobiotic piglets by *Helicobacter pylori* deficient in two flagellin genes. *Infect. Immun.* **64**:2445–2448.
41. Exner, M., P. Doig, T. Trust, and R. Hancock. 1995. Isolation and characterization of a family of porin proteins from *Helicobacter pylori*. *Infect. Immun.* **63**:1567–1572.
42. Forsyth, M. H., J. C. Atherton, M. J. Blaser, and T. L. Cover. 1998. Heterogeneity in levels of vacuolating cytotoxin gene (*vacA*) transcription among *Helicobacter pylori* strains. *Infect. Immun.* **66**:3088–3094.
43. Fraser, C. M., S. J. Norris, G. M. Weinstock, O. White, G. G. Sutton, R. Dodson, M. Gwinn, E. K. Hickey, R. Clayton, K. A. Ketchum, E. Sodergren, J. M. Hardham, M. P. McLeod, S. Salzberg, J. Peterson, H. Khalak, D. Richardson, J. K. Howell, M. Chidambaram, T. Utterback, L. McDonald, P. Artiach, C. Bowman, M. D. Cotton, J. C. Venter, et al. 1998. Complete genome sequence of *Treponema pallidum*, the syphilis spirochete. *Science* **281**:375–388.
44. Fredrick, K., and J. Helmann. 1994. Dual chemotaxis signaling pathways in *Bacillus subtilis*: a sigma D-dependent gene encodes a novel protein with both CheW and CheY homologous domains. *J. Bacteriol.* **176**:2727–2735.
45. Garner, R. M., J. J. Fulkerson, and H. L. Mobley. 1998. *Helicobacter pylori* glutamine synthetase lacks features associated with transcriptional and posttranslational regulation. *Infect. Immun.* **66**:1839–1847.
46. Ge, Z., Q. Jiang, M. S. Kalisiak, and D. E. Taylor. 1997. Cloning and functional characterization of *Helicobacter pylori* fumarate reductase operon comprising three structural genes coding for subunits C, A and B. *Gene* **204**:227–234.
47. Ge, Z., and D. E. Taylor. 1996. *Helicobacter pylori* genes *hpcopA* and *hpcopP* constitute a cop operon involved in copper export. *FEMS Microbiol. Lett.* **145**:181–188.
48. Ge, Z., and D. E. Taylor. 1996. Sequencing, expression, and genetic characterization of the *Helicobacter pylori* *ftsH* gene encoding a protein homologous to members of a novel putative ATPase family. *J. Bacteriol.* **178**:6151–6157.
49. Ghiara, P., M. Marchetti, M. J. Blaser, M. K. Tummuru, T. L. Cover, E. D. Segal, L. S. Tompkins, and R. Rappuoli. 1995. Role of the *Helicobacter pylori* virulence factors vacuolating cytotoxin, CagA, and urease in a mouse model of disease. *Infect. Immun.* **63**:4154–4160.
50. Ghuysen, J.-M. 1994. Serine  $\beta$ -lactamases and penicillin-binding proteins. *Annu. Rev. Microbiol.* **45**:37–67.
51. Goldman, B. S., and R. G. Kranz. 1998. Evolution and horizontal transfer of an entire biosynthetic pathway for cytochrome C biogenesis—*Helicobacter*, *Deinococcus*, *Archaea* and more. *Mol. Microbiol.* **27**:871–873.
52. Goodwin, C. S., M. D. Collins, and E. Blincow. 1986. The absence of thermoplasmaquinones in *Campylobacter pyloridis*, and its temperature and pH growth range. *Microbiol. Lett.* **32**:137–140.
53. Hale, C. A., and P. A. J. deBoer. 1997. Direct binding of FtsZ to ZipA, an essential component of the septal ring structure that mediates cell division in *Escherichia coli*. *Cell* **96**:1–20.
54. Han, J., E. Yu, I. Lee, and Y. Lee. 1997. Diversity among clinical isolates of *Helicobacter pylori* in Korea. *Mol. Cells* **7**:544–547.
55. Hancock, R. E., K. Hantke, and V. Braun. 1997. Iron transport in *Escherichia coli* K-12. 2,3-Dihydroxybenzoate-promoted iron uptake. *Arch. Microbiol.* **114**:231–239.
56. Haque, M., Y. Hirai, K. Yokota, N. Mori, I. Jahan, H. Ito, H. Hotta, I. Yano, Y. Kanemasa, and K. Oguma. 1996. Lipid profile of *Helicobacter* spp.: presence of cholesterol glucoside as a characteristic feature. *J. Bacteriol.* **178**:2065–2070.
57. Hazell, S. L., D. J. Evans, Jr., and D. Y. Graham. 1991. *Helicobacter pylori* catalase. *J. Gen. Microbiol.* **137**:57–61.
58. Hirai, Y., M. Haque, T. Yoshida, K. Yokota, T. Yasuda, and K. Oguma. 1995. Unique cholesteryl glucosides in *Helicobacter pylori*: composition and structural analysis. *J. Bacteriol.* **177**:5327–5333.
59. Hoffman, P. S., A. Goodwin, J. Johnsen, K. Magee, and S. J. Veldhuyzen van Zanten. 1996. Metabolic activities of metronidazole-sensitive and -resistant strains of *Helicobacter pylori*: repression of pyruvate oxidoreductase and expression of isocitrate lyase activity correlate with resistance. *J. Bacteriol.* **178**:4822–4829.
60. Hofreuter, D., S. Odenbreit, G. Henke, and R. Haas. 1998. Natural competence for DNA transformation in *Helicobacter pylori*: identification and genetic characterization of the *comB* locus. *Mol. Microbiol.* **28**:1027–1038.
61. Holt, J. G., N. R. Krieg, P. H. A. Sneath, J. T. Staley, and S. T. Williams. 1994. Bergey's manual of determinative bacteriology, 9th ed. The Williams & Wilkins Co., Baltimore, Md.
62. Holtje, J. V. 1998. Growth of the stress-bearing and shape-maintaining murine sacculus of *Escherichia coli*. *Microbiol. Mol. Biol. Rev.* **62**:181–203.
63. Hughes, N. J., P. A. Chalk, C. L. Clayton, and D. J. Kelly. 1995. Identification of carboxylation enzymes and characterization of a novel four-subunit pyruvate:flavodoxin oxidoreductase from *Helicobacter pylori*. *J. Bacteriol.* **177**:3953–3959.
64. Hughes, N. J., C. L. Clayton, P. A. Chalk, and D. J. Kelly. 1998. *Helicobacter pylori* *porCDAB* and *oorDABC* genes encode distinct pyruvate:flavodoxin and 2-oxoglutarate:acceptor oxidoreductases which mediate electron transport to NADP. *J. Bacteriol.* **180**:1119–1128.
65. Hunt, R. 1996. The role of *Helicobacter pylori* in pathogenesis: the spectrum of clinical outcomes. *Scand. J. Gastroenterol. Suppl.* **220**:3–9.
66. Hurtado, A., R. J. Owen, and M. Desai. 1994. Flagellin gene profiling of *Helicobacter pylori* infecting symptomatic and asymptomatic individuals. *Res. Microbiol.* **145**:585–594.
67. Husson, M. O., D. Legrand, G. Spik, and H. Leclerc. 1993. Iron acquisition by *Helicobacter pylori*: importance of human lactoferrin. *Infect. Immun.* **61**:2694–2697.
68. Ikeda, F., Y. Yokota, Y. Mine, and M. Tatsuta. 1990. Activity of cefixime against *Helicobacter pylori* and affinities for the penicillin-binding proteins. *Antimicrob. Agents Chemother.* **34**:2426–2428.
69. Ikeda, M., T. Sato, M. Wachi, H. K. Jung, F. Ishino, Y. Kobayashi, and M. Matsushashi. 1989. Structural similarity among *Escherichia coli* FtsW and RodA proteins and *Bacillus subtilis* SpoVE protein, which function in cell division, cell elongation, and spore formation, respectively. *J. Bacteriol.* **171**:6375–6378.
70. Illingworth, D. S., K. S. Walter, P. L. Griffiths, and R. Barclay. 1993. Siderophore production and iron-regulated envelope proteins of *Helicobacter pylori*. *Int. J. Med. Microbiol. Virol. Parasitol. Infect. Dis.* **280**:113–119.
71. Iver, D., A. Arnqvist, I.-M. Frick, L. Engstrand, A. Covacci, K.-E. Magnusson, and T. Boren. 1996. The *Helicobacter pylori* blood group antigen binding adhesin. *Gut* **39**:A55. (Abstract.)
72. Iver, D., A. Arnqvist, J. Ogren, I. Frick, D. Kersulyte, E. Incecik, D. Berg, A. Covacci, L. Engstrand, and T. Boren. 1998. *Helicobacter pylori* adhesin binding fucosylated histo-blood group antigens revealed by retagging. *Science* **279**:373–377.
73. Inamoto, Y., S. Ariyama, Y. Hamanaka, K. Okita, Y. Kaneta, T. Nagate, I. Kondou, and T. Takemoto. 1993. Lipid analysis of *Helicobacter pylori*. *J. Clin. Gastroenterol.* **17**:S136–S139.
74. Ishino, F., W. Park, S. Tomioka, S. Tamaki, and I. Takase. 1986. Peptidoglycan synthetic activities in membranes of *Escherichia coli* caused by overproduction of penicillin-binding protein 2 and RodA protein. *J. Biol. Chem.* **261**:7024–7031.
75. Jiang, Q., K. Hiratsuka, and D. Taylor. 1996. Variability of gene order in different *Helicobacter pylori* strains contributes to genome diversity. *Mol. Microbiol.* **20**:833–842.
76. Kadner, R. 1996. Cytoplasmic membrane. p. 58–87. *In* F. C. Neidhardt, R. Curtiss III, J. L. Ingraham, E. C. C. Lin, K. B. Low, B. Magasanik, W. S. Reznikoff, M. Riley, M. Schaechter, and H. E. Umberger (ed.), *Escherichia coli* and *Salmonella*: cellular and molecular biology, vol. 1. ASM Press, Washington, D.C.
77. Kansau, I., J. Raymond, E. Bingen, P. Courcoux, N. Kalach, M. Bergeret, N. Braimi, C. Dupont and A. Labigne. 1996. Genotyping of *Helicobacter pylori* isolates by sequencing of PCR products and comparison with the RAPD technique. *Res. Microbiol.* **147**:661–669.
78. Karita, M., M. Tsuda, and T. Nakazawa. 1995. Essential role of urease *in vitro* and *in vivo* *Helicobacter pylori* colonization study using a wild-type and isogenic urease mutant strain. *J. Clin. Gastroenterol.* **21**:S160–163.
79. Kersulyte, D., N. S. Akopyants, S. Clifton, B. S. Roe, and D. E. Berg. 1998. Novel sequence organization and insertion specificity of IS605 and IS606: chimeric transposable elements of *Helicobacter pylori*. *Gene* **223**:175–186.
80. Klebba, P. E., J. M. Rutz, J. Liu, and C. K. Murphy. 1993. Mechanisms of TonB-catalyzed iron transport through the enteric bacterial cell envelope. *J. Bioenerg. Biomembr.* **25**:603–611.
81. Klenk, H. P., R. A. Clayton, J. F. Tomb, O. White, K. E. Nelson, K. A. Ketchum, R. J. Dodson, M. Gwinn, E. K. Hickey, J. D. Peterson, D. L. Richardson, A. R. Kerlavage, D. E. Graham, N. C. Kyrpides, R. D. Fleischmann, J. Quackenbush, N. H. Lee, G. G. Sutton, S. Gill, E. F. Kirkness, B. A. Dougherty, K. McKenney, M. D. Adams, B. Loftus, J. C. Venter, et al. 1997. The complete genome sequence of the hyperthermophilic, sulphate-reducing archaeon *Archaeoglobus fulgidus*. *Nature* **390**:364–370.
82. Komorowska, M., H. Szafran, T. Popiela, and Z. Szafran. 1981. Free amino acids in gastric juice. *Acta Physiol. Pol.* **32**:559–567.
83. Kostrzynska, M., P. W. O'Toole, D. E. Taylor, and T. J. Trust. 1994.

- Molecular characterization of a conserved 20-kilodalton membrane-associated lipoprotein antigen of *Helicobacter pylori*. *J. Bacteriol.* **176**:5938–5948.
84. Kunst, F., N. Ogasawara, I. Moszer, A. Albertini, G. Alloni, V. Azevedo, M. Bertero, P. Bessieres, A. Bolotin, S. Borchert, R. Borriss, B. Boursier, A. Brans, M. Braun, S. Brignell, S. Bron, S. Brouillet, C. Bruschi, B. Caldwell, V. Capuano, N. Carter, S.-K. Choi, J.-J. Codani, I. Connerton, N. Cummings, R. Daniel, F. Denizot, K. Devine, A. Dusterhoft, S. Ehrlich, P. Emmerson, K. Entian, J. Errington, C. Fabret, E. Ferrari, D. Foulger, C. Fritz, M. Fujita, Y. Fujita, S. Fuma, A. Galizzi, N. Galleron, S.-Y. Ghim, P. Glaser, A. Goffeau, E. Golightly, G. Grandi, G. Guiseppi, B. Guy, K. Haga, J. Haiech, C. Harwood, A. Henaut, H. Hilbert, S. Holsappel, S. Hosono, M.-F. Hullo, M. Itaya, L. Jones, B. Joris, D. Karamata, Y. Kasahara, M. Klaerr-Blanchard, C. Klein, Y. Kobayashi, P. Loetter, G. Koningsstein, S. Krogh, M. Kumano, K. Kurita, A. Lapidus, S. Lardinois, J. Lauber, V. Lazarevic, S.-M. Lee, A. Levine, H. Liu, S. Masuda, C. Mauel, C. Medigue, N. Median, R. Mellado, M. Mizuno, D. Moestl, S. Nakai, M. Noback, D. Noone, M. O'Reilly, K. Ogawa, A. Ogiwara, B. Oudega, S.-H. Park, V. Parro, T. Pohl, D. Portetelle, S. Porwollik, A. Prescott, E. Presecan, P. Pujic, B. Purnelle, et al. 1997. The complete genome sequence of the Gram-positive bacterium *Bacillus subtilis*. *Nature* **390**:249–256.
  85. Kusukawa, N., T. Yura, C. Ueguchi, Y. Akiyama, and K. Ito. 1989. Effects of mutations in heat-shock genes *groES* and *groEL* on protein export in *Escherichia coli*. *EMBO J.* **8**:3517–3521.
  86. Labigne, A., V. Cussac, and P. Courcoux. 1991. Shuttle cloning and nucleotide sequences of *Helicobacter pylori* genes responsible for urease activity. *J. Bacteriol.* **173**:1920–1931.
  87. Labigne, A., and H. deReuse. 1996. Determinants of *Helicobacter pylori* pathogenicity. *Infect. Agents Dis.* **5**:191–202.
  88. Lee, C. A. 1997. Type III secretion systems: machines to deliver bacterial proteins into eukaryotic cells? *Trends Microbiol.* **5**:148–156.
  89. Leying, H., S. Suerbaum, G. Geis, and R. Haas. 1992. Cloning and genetic characterization of a *Helicobacter pylori* flagellin gene. *Mol. Microbiol.* **6**:2863–2874.
  90. Luke, C., and C. Penn. 1995. Identification of a 29kDa flagellar sheath protein in *Helicobacter pylori* using a murine monoclonal antibody. *Microbiology* **141**:597–604.
  91. Luke, C. J., E. Kubiak, A. Cockayne, T. S. Elliott, and C. W. Penn. 1990. Identification of flagellar and associated polypeptides of *Helicobacter* (formerly *Campylobacter*) *pylori*. *FEMS Microbiol. Lett.* **59**:225–230.
  92. Mahdi, A. A., and R. G. Lloyd. 1989. Identification of the *recR* locus of *Escherichia coli* K-12 and analysis of its role in recombination and DNA repair. *Mol. Gen. Genet.* **216**:503–510.
  93. Maier, R. J., C. Fu, J. Gilbert, F. Moshiri, J. Olson, and A. G. Plaut. 1996. Hydrogen uptake hydrogenase in *Helicobacter pylori*. *FEMS Microbiol. Lett.* **141**:71–76.
  94. Marcelli, S. W., H. T. Chang, T. Chapman, P. A. Chalk, R. J. Miles, and R. K. Poole. 1996. The respiratory chain of *Helicobacter pylori*: identification of cytochromes and the effects of oxygen on cytochrome and menaquinone levels. *FEMS Microbiol. Lett.* **138**:59–64.
  95. Marshall, D. G., W. G. Dundon, S. M. Beesley, and C. J. Smyth. 1998. *Helicobacter pylori*—a conundrum of genetic diversity. *Microbiology* **144**:2925–2939.
  96. Martin, S., M. Edbrooke, T. Hodgman, D. van der Eijnden, and M. Bird. 1997. Lewis X biosynthesis in *Helicobacter pylori*. Molecular cloning of an  $\alpha(1,3)$ -fucosyltransferase gene. *J. Biol. Chem.* **272**:21349–21356.
  97. Matin, A., E. Zychlinsky, M. Keyhan, and G. Sachs. 1996. Capacity of *Helicobacter pylori* to generate ionic gradients at low pH is similar to that of bacteria which grow under strongly acidic conditions. *Infect. Immun.* **64**:1434–1436.
  98. Mendz, G. L., B. P. Burns, and S. L. Hazell. 1995. Characterisation of glucose transport in *Helicobacter pylori*. *Biochim. Biophys. Acta* **1244**:269–276.
  99. Mendz, G. L., and S. L. Hazell. 1995. Amino acid utilization by *Helicobacter pylori*. *Int. J. Biochem. Cell Biol.* **27**:1085–1093.
  100. Mendz, G. L., and S. L. Hazell. 1991. Evidence for a pentose phosphate pathway in *Helicobacter pylori*. *FEMS Lett.* **84**:331–336.
  101. Mendz, G. L., and S. L. Hazell. 1993. Fumarate catabolism in *Helicobacter pylori*. *Biochem. Mol. Biol. Int.* **31**:325–332.
  102. Mendz, G. L., and S. L. Hazell. 1993. Glucose phosphorylation in *Helicobacter pylori*. *Arch. Biochem. Biophys.* **300**:522–525.
  103. Mendz, G. L., S. L. Hazell, and B. P. Burns. 1994. The Entner-Doudoroff pathway in *Helicobacter pylori*. *Arch. Biochem. Biophys.* **312**:349–356.
  104. Mendz, G. L., S. L. Hazell, and B. P. Burns. 1993. Glucose utilization and lactate production by *Helicobacter pylori*. *J. Gen. Microbiol.* **139**:3023–3028.
  105. Mendz, G. L., S. L. Hazell, and S. Srinivasan. 1995. Fumarate reductase: a target for therapeutic intervention against *Helicobacter pylori*. *Arch. Biochem. Biophys.* **321**:153–159.
  106. Mendz, G. L., S. L. Hazell, and L. van Gorkom. 1994. Pyruvate metabolism in *Helicobacter pylori*. *Arch. Microbiol.* **162**:187–192.
  107. Mendz, G. L., B. M. Jimenez, S. L. Hazell, A. M. Gero, and W. J. O'Sullivan. 1994. Salvage synthesis of purine nucleotides by *Helicobacter pylori*. *J. Appl. Bacteriol.* **77**:674–681.
  108. Mendz, G. L., B. M. Jimenez, S. L. Hazell, A. M. Gero, and W. J. O'Sullivan. 1994. De novo synthesis of pyrimidine nucleotides by *Helicobacter pylori*. *J. Appl. Bacteriol.* **77**:1–8.
  109. Miehke, S., A. Meining, A. Morgner, E. Bayerdorffer, N. Lehn, M. Stolte, D. Y. Graham, and M. F. Go. 1998. Frequency of *vacA* genotypes and cytotoxin activity in *Helicobacter pylori* associated with low-grade gastric mucosa-associated lymphoid tissue lymphoma. *J. Clin. Microbiol.* **36**:2369–2370.
  110. Mizote, T., H. Yoshizawa, and T. Nakazawa. 1997. Urease-independent chemotactic responses of *Helicobacter pylori* to urea, urease inhibitors, and sodium bicarbonate. *Infect. Immun.* **65**:1519–1521.
  111. Mobley, H. L. 1996. Defining *Helicobacter pylori* as a pathogen: strain heterogeneity and virulence. *Am. J. Med.* **100**:2S–9S.
  112. Mobley, H. L. 1997. *Helicobacter pylori* factors associated with disease development. *Gastroenterology* **113**:S21–S28.
  113. Mobley, H. L., M. D. Island, and R. P. Hausinger. 1995. Molecular biology of microbial ureases. *Microbiol. Rev.* **59**:451–480.
  114. Mobley, H. L. T., R. M. Garner, and P. Bauerfeind. 1995. *Helicobacter pylori* nickel-transport gene *nixA*: synthesis of catalytically active urease in *Escherichia coli* independent of growth conditions. *Mol. Microbiol.* **16**:97–109.
  115. Moran, A. P., B. Lindner, and E. Walsh. 1997. Structural characterization of the lipid A component of *Helicobacter pylori* rough- and smooth-form lipopolysaccharides. *J. Bacteriol.* **179**:6453–6463.
  116. Nagata, K., S. Tsukita, T. Tamura, and N. Sone. 1996. A cb-type cytochrome-c oxidase terminates the respiratory chain in *Helicobacter pylori*. *Microbiology* **142**:1757–1763.
  117. Nakao, H., S. Shioda, and S. Yamamoto. 1990. Purification and properties of carboxynorspermidine decarboxylase, a novel enzyme involved in norspermidine biosynthesis from *Vibrio alginolyticus*. *J. Gen. Microbiol.* **136**:1699–1704.
  118. Nedeskov, P. 1994. Nutritional requirements for growth of *Helicobacter pylori*. *Appl. Environ. Microbiol.* **60**:3450–3453.
  119. Neidhardt, F. C., R. Curtiss III, J. L. Ingraham, E. C. C. Lin, K. B. Low, B. Magasanik, W. S. Reznikoff, M. Riley, M. Schaechter, and H. E. Umberger (ed.). 1996. *Escherichia coli* and *Salmonella*: cellular and molecular biology, 2nd ed. ASM Press, Washington, D.C.
  120. Newell, D. 1987. Identification of the outer membrane proteins of *Campylobacter pyloridis* and antigenic cross-reactivity between *C. pyloridis* and *C. jejuni*. *J. Gen. Microbiol.* **133**:163–170.
  121. Nishiyama, K., S. Mizushima, and H. Tokuda. 1993. A novel membrane protein involved in protein translocation across the cytoplasmic membrane of *Escherichia coli*. *EMBO J.* **12**:3409–3415.
  122. Odenbreit, S., B. Wieland, and R. Haas. 1996. Cloning and genetic characterization of *Helicobacter pylori* catalase and construction of a catalase-deficient mutant strain. *J. Bacteriol.* **178**:6960–6967.
  123. Ohta-Tada, U., A. Takagi, Y. Koga, S. Kamiya, and T. Miwa. 1997. Flagellin gene diversity among *Helicobacter pylori* strains and IL-8 secretion from gastric epithelial cells. *Scand. J. Gastroenterol.* **32**:455–459.
  124. O'Toole, P. W., M. Kostrzynska, and T. J. Trust. 1994. Non-motile mutants of *Helicobacter pylori* and *Helicobacter mustelae* defective in flagellar hook production. *Mol. Microbiol.* **14**:691–703.
  125. Page, M. D., E. Tomlinson, and S. J. Ferguson. 1997. Unexpected implications from the *Helicobacter pylori* genome for understanding periplasmic c-type cytochrome assembly in gram-negative bacteria in coexistence with disulphide bond formation. *Mol. Microbiol.* **26**:413–415.
  126. Pahel, G., A. D. Zelenetz, and B. M. Tyler. 1978. *glbB* gene and regulation of nitrogen metabolism by glutamine synthetase in *Escherichia coli*. *J. Bacteriol.* **133**:139–148.
  127. Park, J. T. 1995. Why does *Escherichia coli* recycle its cell wall peptides? *Mol. Microbiol.* **17**:421–426.
  128. Penn, C. W., and C. J. Luke. 1992. Bacterial flagellar diversity and significance in pathogenesis. *FEMS Microbiol. Lett.* **79**:331–336.
  129. Phadnis, S. H., M. H. Parlow, M. Levy, D. Iver, C. M. Caulkins, J. B. Connors, and B. E. Dunn. 1996. Surface localization of *Helicobacter pylori* urease and a heat shock protein homolog requires bacterial autolysis. *Infect. Immun.* **64**:905–912.
  130. Preisig, O., D. Anthamatten, and H. Henneke. 1993. Genes for a microaerobically induced oxidase complex in *Bradyrhizobium japonicum* are essential for a nitrogen-fixing endosymbiosis. *Proc. Nat. Acad. Sci. USA* **90**:3309–3313.
  131. Reitzer, L. 1996. Ammonia assimilation and the biosynthesis of glutamine, glutamate, aspartate, asparagine, L-alanine, and D-alanine, p. 391–407. *In* F. C. Neidhardt, R. Curtiss III, J. L. Ingraham, E. C. C. Lin, K. B. Low, B. Magasanik, W. S. Reznikoff, M. Riley, M. Schaechter, and H. E. Umberger (ed.), *Escherichia coli* and *Salmonella*: cellular and molecular biology, vol. 1. ASM Press, Washington, D.C.
  132. Reynolds, D. J., and C. W. Penn. 1994. Characteristics of *Helicobacter pylori* growth in a defined medium and determination of its amino acid requirements. *Microbiology* **140**:2649–2656.
  133. Roine, R. P., K. S. Salmela, and M. Salaspuuro. 1995. Alcohol metabolism in *Helicobacter pylori*-infected stomach. *Ann. Med.* **27**:583–588.
  134. Rosario, M. M., K. L. Fredrick, G. W. Ordal, and J. D. Helmann. 1994.

- Chemotaxis in *Bacillus subtilis* requires either of two functionally redundant CheW homologs. *J. Bacteriol.* **176**:2736–2739.
135. Salmela, K. S., R. P. Roine, J. Hook-Nikanne, T. U. Kosunen, and M. Salaspuro. 1994. Acetaldehyde and ethanol production by *Helicobacter pylori*. *Scand. J. Gastroenterol.* **29**:309–312.
  136. Salmela, K. S., R. P. Roine, T. Koivisto, J. Hook-Nikanne, T. U. Kosunen, and M. Salaspuro. 1993. Characteristics of *Helicobacter pylori* alcohol dehydrogenase. *Gastroenterology* **105**:325–330.
  137. Segal, E. D., C. Lange, A. Covacci, L. S. Tompkins, and S. Falkow. 1997. Induction of host signal transduction pathways by *Helicobacter pylori*. *Proc. Natl. Acad. Sci. USA* **94**:7595–7599.
  138. Seifert, J., N. Kunz, R. Flachmann, A. Laufer, K. D. Jany, and H. G. Gassen. 1990. Expression of the *E. coli nadB* gene and characterization of the gene product L-aspartate oxidase. *Biol. Chem. Hoppe-Seyley* **371**:239–248.
  139. Smith, D., L. Douchette-Stamm, C. Deloughery, H. Lee, J. Dubois, T. Aldredge, R. Bashirzadeh, D. Blakely, R. Cook, K. Gilbert, D. Harrison, L. Hoang, P. Keagle, W. Lumm, B. Pothier, D. Qiu, R. Spadafors, R. Vicaire, Y. Wang, J. Wierzbowski, R. Gibson, N. Jiwani, A. Caruso, D. Bush, H. Safer, D. Patwell, S. Prabhakar, S. McDougall, G. Shimer, A. Goyal, S. Pietrokovski, G. Church, C. Daniels, J.-I. Mao, P. Rice, J. Nolling, and J. Reeve. 1997. Complete genome sequence of *Methanobacterium thermoautotrophicum* deltaH: functional analysis and comparative genomics. *J. Bacteriol.* **179**:7135–7155.
  140. Spiegelhalder, C., B. Gerstenecker, A. Kersten, E. Schiltz, and M. Kist. 1993. Purification of *Helicobacter pylori* superoxide dismutase and cloning and sequencing of the gene. *Infect. Immun.* **61**:5315–5325.
  141. Spohn, G., D. Beier, R. Rappuoli, and V. Scarlato. 1997. Transcriptional analysis of the divergent *cagAB* genes encoded by the pathogenicity island of *Helicobacter pylori*. *Mol. Microbiol.* **26**:361–372.
  142. Spratt, B. G. 1977. Properties of the penicillin-binding proteins of *Escherichia coli* K12. *Eur. J. Biochem.* **72**:341–352.
  143. Stark, R. M., M. S. Suleiman, I. J. Hassan, J. Greenman, and M. R. Millar. 1997. Amino acid utilisation and deamination of glutamine and asparagine by *Helicobacter pylori*. *J. Med. Microbiol.* **46**:793–800.
  144. Stephens, J. C., J. A. Stewart, A. M. Folwell, and B. J. Rathbone. 1998. *Helicobacter pylori cagA* status, *vacA* genotypes and ulcer disease. *Eur. J. Gastroenterol. Hepatol.* **10**:381–384.
  145. Stock, A., E. Schaeffer, D. E. Koshland, and J. Stock. 1987. A second type of protein methylation reaction in bacterial chemotaxis. *J. Biol. Chem.* **262**:8011–8014.
  146. Strauch, M. A., H. Zalkin, and A. I. Aronson. 1988. Characterization of the glutamyl-tRNA(Gln)-to-glutamyl-tRNA(Gln) amidotransferase reaction of *Bacillus subtilis*. *J. Bacteriol.* **170**:916–920.
  147. Suerbaum, S., C. Josenhans, and A. Labigne. 1993. Cloning and genetic characterization of the *Helicobacter pylori* and *Helicobacter mustellae flaB* genes and construction of *H. pylori flaA*- and *flaB*-negative mutants by allelic exchange. *J. Bacteriol.* **175**:3278–3288.
  148. Taylor, D., M. Eaton, N. Chang, and S. Salama. 1992. Construction of a *Helicobacter pylori* genome map and demonstration of diversity at the genome level. *J. Bacteriol.* **174**:6800–6806.
  149. Tomb, J.-F., O. White, A. Kerlavage, R. Clayton, G. Sutton, R. Fleischmann, K. Ketchum, H. Klenk, S. Gill, B. Dougherty, K. Nelson, J. Quackenbush, L. Zhou, E. Kirkness, S. Peterson, B. Loftus, D. Richardson, R. Dodson, H. Khalak, A. Glodek, K. McKenney, L. Fitzgerald, N. Lee, M. Adams, E. Hickey, D. Berg, J. Gocayne, T. Utterback, J. Peterson, J. Kelley, M. Cotton, J. Weidman, C. Fujii, C. Bowman, L. Wathley, E. Wallin, W. Hayes, M. Borodovsky, P. Karp, H. Smith, C. Fraser, and J. Venter. 1997. The complete genome sequence of the gastric pathogen *Helicobacter pylori*. *Nature* **388**:539–547.
  150. Tsuda, M., M. Karita, M. Morshed, K. Okita, and T. Nakazawa. 1994. A urease-negative mutant of *Helicobacter pylori* constructed by allelic exchange mutagenesis lacks the ability to colonize the nude mouse stomach. *Infect. Immun.* **62**:3586–3589.
  151. van Doorn, L. J., C. Figueiredo, R. Sanna, S. Pena, P. Midolo, E. K. Ng, J. C. Atherton, M. J. Blaser, and W. G. Quint. 1998. Expanding allelic diversity of *Helicobacter pylori vacA*. *J. Clin. Microbiol.* **36**:2597–2603.
  152. van Doorn, L. J., C. Figueiredo, R. Sanna, A. Plaisier, P. Schneeberger, W. de Boer, and W. Quint. 1998. Clinical relevance of the *cagA*, *vacA*, and *iceA* status of *Helicobacter pylori*. *Gastroenterology* **115**:58–66.
  153. Vanet, A., and A. Labigne. 1998. Evidence for specific secretion rather than autolysis in the release of some *Helicobacter pylori* proteins. *Infect. Immun.* **66**:1023–1027.
  154. Vincente, M., M. J. Gomez, and J. A. Agata. 1990. Regulation of transcription of cell division genes in the *Escherichia coli dcw* cluster. *Cell. Mol. Life Sci.* **59**:317–324.
  155. Weiss, A. A., F. D. Johnson, and D. L. Burns. 1993. Molecular characterization of an operon required for pertussis toxin secretion. *Proc. Natl. Acad. Sci. USA* **90**:2970–2974.
  156. Westblom, T. U., S. Phadnis, W. Langenberg, K. Yoneda, E. Madan, and B. R. Midkiff. 1992. Catalase negative mutants of *Helicobacter pylori*. *Eur. J. Clin. Microbiol. Infect. Dis.* **11**:522–526.
  157. Williams, C. L., T. Preston, M. Hossack, C. Slater, and K. E. McColl. 1996. *Helicobacter pylori* utilises urea for amino acid synthesis. *FEMS Immunol. Med. Microbiol.* **13**:87–94.
  158. Worst, D. J., M. M. Gerrits, C. M. Vandenbroucke-Grauls, and J. G. Kusters. 1998. *Helicobacter pylori ribBA*-mediated riboflavin production is involved in iron acquisition. *J. Bacteriol.* **180**:1473–1479.
  159. Xu, Q., R. M. J. Peek, G. G. Miller, and M. J. Blaser. 1997. The *Helicobacter pylori* genome is modified at CATG by the product of *hpyIM*. *J. Bacteriol.* **179**:6807–6815.
  160. Zakharova, N., P. S. Hoffman, D. E. Berg, and K. Severinov. 1998. The largest subunits of RNA polymerase from gastric helicobacters are tethered. *J. Biol. Chem.* **273**:19371–19374.